



Long-sleeved shirt and long pants,
Shoes plus socks,
Applicators using hand held equipment must also wear chemical-resistant gloves made out of: barrier laminate, butyl rubber >14 mils, nitrile rubber >14 mils, or neoprene rubber >14 mils.
Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

- Keep children and pets off treated area until product has been watered in and lawn is dry.
- Do not apply this product to edible growing crops or stored raw agricultural commodities used for food or feed.
- Do not allow applications to contact water inhabited by fish, such as in ornamental fish ponds that are located around structures being treated.
- Do not apply this product by aerial application.
- Use of this pesticide adjacent to water may affect aquatic organisms. To protect these organisms, do not apply this pesticide within 25 ft. of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.
- In New York State, this product may not be applied within 100 feet of a coastal marsh, or streams that drains directly into a coastal marsh.
- All restrictions and precautions which appear on the label of this product must be followed.
- Do not apply more than 4.6 lb of this product per 1,000 square feet or 200 lb of product (0.4 lb of imidacloprid active ingredient) per acre per year.
- Not for use in commercial greenhouses, nurseries on sod farms or on grass grown for seed

For use on turfgrass in residential lawns, business and office complexes, shopping complexes, multi-family residential complexes, golf courses, airports, cemeteries, parks, playgrounds, and athletic fields to control soil inhabiting pests as listed in the table below. This product can also be used for the suppression of cutworms in turfgrass areas.

Effective control can be achieved when applications are made before or during the egg laying period. The timing of applications can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. For best control apply prior to egg hatch of the target pests. Allow rainfall or irrigate sufficiently within 24 hours after application to move the active ingredient through the thatch.

Do not make applications when the grass areas are waterlogged or the soil is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist. The area to be treated must be able to accept vertical penetration of rainfall or irrigation water. Do not mow turf or lawn area until after sufficient irrigation or rainfall has occurred so that uniformity of application will not be affected.

Consult your local State Agricultural Experiment Station or State Extension Turf Specialists for more specific information regarding timing of application.

Apply this product uniformly over the area being treated with granular application equipment. Both drop-type and rotary-type spreaders may be used. Do not use spreaders which apply the material in concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly and accurately.

Use the higher listed rate for longer length of control. Watering (irrigation or rainfall) following application will dissolve the granules and release the insecticide. Uniform coverage is necessary to provide the most effective level of control. Consult your state university or local Cooperative Extension Service office for specific pest control application timing in your area. Start application to turf prior to the establishment of high insect pest populations and significant turf damage. Make reapplications as necessary to keep pest populations under control, using higher rates as pest pressure increases.

Typically, a lawn utilizes 1 to 4 lbs of nitrogen per 1,000 sq ft per growing season. When applied as directed, this product provides .39 to .82 lbs. of nitrogen per 1,000sq ft. Contact your local Extension Service for information specific to your geographic region.

Earthway 2400 Rotary	15	16	17	18
Lely Broadcast	2.5	3	3.4	3.5

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY ANY COMMUNICATION TO EITHER TO CUSTOMERS OR IF ANY, OR FOR ANY DAMAGES OR DAMAGES OF ANY KIND OR AMOUNT, DEMANDS FOR RECALL, RECALL COSTS, OR FOR ANY LOSS OF PROFITS, OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(RV082517)



NET WT. 50 lbs. (22.7 kg)

Distributed By
Eau Claire Co-op Oil Co.
Box 837, Eau Claire, WI 54702

Doc# RP228-95-18

Imidacloprid	0.20%
Lambda-cyhalothrin	0.04%
Other Ingredients:	99.76%
TOTAL:	100.00%

EPA Est. No. 57131-WI-1

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

Causes moderate eye irritation. Avoid contact with eyes or clothing.

This product contains a chemical with properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.