



17-0-5 Fertilizer with 0.069% ProSect® Insecticide

- Contains 0.069% ProSect® (bifenthrin), a pyrethroid insecticide
- ProSect® can be used to control a wide variety of insects as listed on the label
- The 17-0-5 fertilizer contains 30% Fortify®-N



PRODUCT CODE ASP175UBF5

DESCRIPTION 17-0-5 / 0.069% bifenthrin / 30% Fortify®-N

PRODUCT SIZE SGN 215

N SOURCE Urea and Fortify®-N

APPLICATION 3.33 to 6.66 lbs of product / per 1,000 ft²

PRODUCT USES Controls a variety of insects including: Annual bluegrass weevil adults, Ants, Armyworms, Billbug adults, Black turfgrass ataeinus adults, Centipedes, Chiggers, Chinch bugs, Cutworms, Fire ants, Fleas, Imported fire ants, Leafhoppers, Mealybugs, Millipedes, Mole crickets, Sod webworms and Ticks. Please see label for rates and directions for use.

PACKAGING 50 lb. bags x 40 per pallet

To order, contact your local distributor from The Andersons
Visit our website: www.andersonsplantnutrient.com
or contact us at **800-253-5296**



17-0-5 Fertilizer

with 0.069% *ProSect*[®] Insecticide

FOR USE BY INDIVIDUALS/FIRMS LICENSED OR REGISTERED BY THE STATE TO APPLY INSECTICIDE PRODUCTS.

FOR USE TO CONTROL ANTS (INCLUDING IMPORTED FIRE ANTS), MOLE CRICKETS AND OTHER LISTED PESTS ON LAWN IN LANDSCAPED AREAS AROUND RESIDENTIAL; INSTITUTIONAL, PUBLIC, COMMERCIAL AND INUDSTRIAL BUILDINGS, PARKS, RECREATIONAL AREAS AND ATHLETIC FIELDS.

ACTIVE INGREDIENT:	
Bifenthrin: *	0.069%
OTHER INGREDIENTS:	99.931%
TOTAL:	100.000%

*Cis isomers 97% min.; trans isomers 3% max.

EPA Reg. No. 5481-539-9198
EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A
Underlined letter is first letter used in run code on bag

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice
EMERGENCY INFORMATION	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-264-3910 or Chemtrec; 1-800-424-9300 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
This product is a pyrethroid. If large amounts have been ingested, milk, cream and other digestible fats and oils may increase absorption and so should be avoided.	

PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

Additional Precautions when Used in the State of New York:

In New York State this product should not be applied to any grass or turfgrass within 100 feet of a water body (lake, pond, river, stream, wetland, drainage ditch). In New York State, make only a single repeat application if there are signs of renewed insect or pest activity, but no sooner than 2 weeks after the first application.

Guaranteed Analysis

Total Nitrogen (N)	17.00%
17.00% Urea Nitrogen (N)*	
Soluble Potash (K ₂ O)	5.00%
Plant nutrients derived from urea and potassium chloride.	
Chlorine (Cl) Max	3.86%
* 5.10% nitrogen stabilized with with N-(n-butyl) thiophosphoric triamide.	

F002091

FLORIDA APPLICATIONS:

We recommend that you follow the Florida Green Industries BMP's at http://www.flaes.org/pdf/BMP_Book_final.pdf

We recommend that you follow the Florida Golf Course BMP's at <http://www.flaes.org/pdf/glfbmp07.pdf>

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production or for research purposes.

Not for use on golf courses, sod farms, nurseries, commercial greenhouses, or grass grown for seed.

Do not apply this product within 25 ft. of lakes, reservoirs, rivers, permanent streams, marshes, natural ponds or commercial fish ponds.

Best results are obtained if the areas are thoroughly watered soon after application.

Do not allow children or pets in treated areas until surfaces are dry.

Do not exceed 13.3 lbs. of product per 1,000 sq. ft. in any given year (580 lbs/acre).

Do not apply by air.

For best results, commence treatment when the pests (listed below) first appear.

Lawn & Perimeter Applications:

Perimeter application: Apply to lawn (or turf) area in a uniform band 6 to 10 ft. wide around and adjacent to structures to help prevent entry of insects into homes or buildings. Use at the rate of 3.33 lb. to 6.66 lb. per 1,000 sq. ft. depending on pest to be treated.

Lawn (or turf) application: Broadcast this product with suitable application equipment which is properly calibrated to ensure uniform coverage over the treatment area. Best results are obtained when two applications at half rates are made perpendicular to each other. Apply 3.33 lb. to 6.66 lb. per 1,000 sq. ft. to control the following pests:

Pest List and Use Rates:

- To control Annual bluegrass weevil adults (*Hyperodes sp.*), Armyworms, Billbug adults, Black turfgrass ataenus adults, Cutworms, Mealybugs and sod webworms, use at the rate of 3.33 lbs/1,000 sq. ft.
- To control Ants, Centipedes, Chiggers (Red bugs), Chinch bugs, Fire ants, Fleas, Imported fire ants, Leafhoppers, Millipedes, Mole crickets and Ticks, use at the rate of 6.66 lbs/1,000 sq. ft.

Note: To improve overall control of Imported fire ant colonies, combine broadcast treatments with appropriate products approved for fire ant mound applications.

READ ENTIRE LABEL BEFORE USING

Spreader Suggestions:

The spreader settings indicated below are approximate and should only be used as a guide. Age and conditions of spreader (especially if parts are worn out) and walking speed can cause variation in application rate. Prior to each use, equipment calibration is recommended. Close spreader hopper and adjust the setting using the suggested settings chart below. If not indicated below, carefully follow spreader manufacturer's recommendations to obtain accurate spread rates. Always fill Spreader on a hard surface to eliminate the chance of spillage on the lawn and close spreader hopper when stopping or turning.

Suggested Spreader Settings				
These suggested spreader settings are not intended to replace calibration. Please calibrate your spreader prior to applying product.				
50 lbs. treats 15,000 sq. ft. (0.34 acre) at 145 lbs. product/acre - LOW RATE (3.33 lbs/1,000 sq. ft.)				
50 lbs. treats 7,500 sq. ft. (0.17 acre) at 290 lbs. product/acre - HIGH RATE (6.66 lbs/1,000 sq. ft.)				
SPREADER	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS	
			LOW RATE 0.10 lb. ai/A	HIGH RATE 0.20 lb. ai/A
Andersons Model 2000 / AccuPro 2000 (cone 4); Andersons 2000 SR / SR-2000 (cone 4)	3 mph	9 feet	L 1/4	P
Andersons LCO-1000	3 mph	9 feet	L 1/4	P
Scotts R-8A (cone 4)	3 mph	9 feet	K 1/4	O
Lesco 020093 (pattern 0.25)	3 mph	6 feet	E	F 1/2
Lesco 705698 (pattern 0.25)	3 mph	6 feet	12	15
Lely Models WTR, WFR, HR, 1250 (PTO at 450 rpm) Pattern II-C	4.5 mph	32 feet	5 1/2	7 1/2
Vicon (03 Series)	4.5 mph	25 feet	27	40

Storage & Disposal

Prohibitions:

Do not contaminate water, food or feed by storage or disposal.

Storage: Store product in its original closed container in a cool, dry place out of reach of children and animals. Carefully open containers and after partial use, close tightly. Avoid excessive heat. If spill, avoid contact. Isolate spills, sweep up, and use as directed on this label.

Pesticide Disposal: Wastes resulting from a spill or from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. If not available, then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. To the extent consistent with applicable law, Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION.

Distributed by:
The Andersons, Inc.
P. O. Box 119
Maumee, OH 43537
www.andersonsplantnutrient.com

Prosect is a registered trademark of The Andersons, Inc.

Net Wt. 50 Lbs. (22.68 kg)



ASP175UBF5