# LebanonTurf

## Country Club®Maximum Dispers Maximum Dispersion

- Utilizes a unique, highly dispersing granule technology
- Contains North Atlantic Ocean sea kelp
- Features METHEX slow release nitrogen
- SGN 80 is ideal for putting greens



#### **GUARANTEED ANALYSIS**

Total Nitrogen (N)	12%	Total Nitrogen (N)		12%
0.9% Ammoniacal Nitrogen		0.9% Ammoniacal Nitrogen		
2.4% Water Insoluble Nitrogen*		2.4% Water Insoluble Nitrog	en*	
4.5% Urea Nitrogen		4.5% Urea Nitrogen		
4.2% Other Water Soluble Nitrogen *		4.2% Other Water Soluble N	itrogen *	
Soluble Potash (K <sub>2</sub> O)				24%
Magnesium (Mg)				
0.5% Water Soluble Magnesium (Mg)	0.070	0.5% Water Soluble Magnes		0.070
Sulfur (S)	0.5%	0.5% Water Soluble Magnes	sum (mg)	0 59/
9.5% Combined Sulfur (S)	5.570	Sultur (S)		9.9%
9.5% Complied Sundi (5)	1 69/	9.5% Combined Sulfur (S)		4.00/
Iron (Fe)	1.0%	Iron (Fe) (actual)		1.6%
0.2% Water Soluble Iron (Fe)		0.2% Water Soluble Iron (Fe	)	
Manganese (Mn)	0.5%	Manganese (Mn)	, 	0.5%
0.5% Water Soluble Manganese (Mn)		0.5% Water Soluble Mangan	nese (Mn)	
Derived From: Ammonium Sulfate, Methylene Ureas, Ur	rea, Sulfate of Potash, Sulfate	Chlorine (Cl) (maximum)		2.0%
of Potash Magnesia, Kelp Meal, Ferric Oxide, Ferrous S		*6.6% Slowly Available Nitrog	gen from Methylene Ureas.	
*6.6% Slowly Available Nitrogen from Methylene Ureas		Derived From: Ammonium Su	ulfate, Methylene Ureas, Urea, Sulfate of	f Potash Sulfate
Chlorine (CI) not more than			al, Ferric Oxide, Ferrous Sulfate, Manga	
ALSO CONTAINS NON PLANT FOOD INGREDIE	ENTS	<b>3</b> / 1	PLANT FOOD INGREDIENTS	
SOIL/PLANT AMENDING INGREDIENTS		SOIL/PLANT AMENDING		
Active Ingredients:		Active Ingredients:		
Humic Acids (derived from Leonardite) 1.09	6		rom Leonardite) 1.0%	
Inert Ingredients (as soil amending) 99.0%			il amending)99.0%	CANAD
99.0% Fertilizer Ingredients		99.0% Fertilizer In		VANAD
romptly after applications by sweeping or blowing. Do not wash ebanon brand fertilizers are formulated for profession		ourses, sports fields, and landsc	apes. Specific application rates and	frequencies are
romptly after applications by sweeping or blowing. Do not wash ebanon brand fertilizers are formulated for professio letermined by the professional applicator based on lo	nal use on turf, including golf c		apes. Specific application rates and	frequencies are
ebanon brand fertilizers are formulated for profession	nal use on turf, including golf c cal soils, regulations, and clim Wa	ate.	Application Rates and Frequencies in Ca	nada: Follow your
ebanon brand fertilizers are formulated for profession	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Wa		Application Rates and Frequencies in Ca professional landscape management pla	nada: Follow your n for your local cours
ebanon brand fertilizers are formulated for profession	nal use on turf, including golf c ical soils, regulations, and clim Hazard Statements Harmful if swallowed.	ate.	Application Rates and Frequencies in Ca professional landscape management pla or area. A suggested application rate is 1	nada: Follow your n for your local cours 1 to 2.8 Kg for 100 m²
ebanon brand fertilizers are formulated for profession	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Wa	ate.	Application Rates and Frequencies in Ca professional landscape management pla	nada: Follow your n for your local cours 1 to 2.8 Kg for 100 m²
ebanon brand fertilizers are formulated for profession	nal use on turf, including golf c ical soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. May be harmful if inhaled.	ate.	Application Rates and Frequencies in Ca professional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc	nada: Follow your n for your local cours 1 to 2.8 Kg for 100 m²
ebanon brand fertilizers are formulated for professio letermined by the professional applicator based on lo Application Rates:	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation.	ate.	Application Rates and Frequencies in Ca professional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc Application Rates:	nada: Follow your n for your local cours 1 to 2.8 Kg for 100 m² :al growing season.
ebanon brand fertilizers are formulated for profession letermined by the professional applicator based on lo Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0)	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation.	arning	Application Rates and Frequencies in Ca professional landscape management pla or area. A suggested application rate is a applied every 4 to 8 weeks during the loc <b>Application Rates:</b> To feed at the rate of 0.24 kg Nitrogen (	nada: Follow your n for your local cours 1 to 2.8 Kg for 100 m <sup>2</sup> al growing season. N) or 0.48 kg
ebanon brand fertilizers are formulated for profession letermined by the professional applicator based on lo Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eve irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling	ate. arning	Application Rates and Frequencies in Ca professional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc Application Rates:	nada: Follow your n for your local cours 1 to 2.8 Kg for 100 m <sup>2</sup> al growing season. N) or 0.48 kg
ebanon brand fertilizers are formulated for profession letermined by the professional applicator based on lo Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlin Wash hands and exposed skin thorough	ate. arning y g nly after handling.	Application Rates and Frequencies in Ca professional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc <b>Application Rates:</b> To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of pr	nada: Follow your n for your local cours t to 2.8 Kg for 100 m <sup>2</sup> cal growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> .
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 103 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 2.1	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlin, Wash hands and exposed skin thorough Do not eat, drink or smoke when using t	ate. Arning J g ny after handling. this product.	Application Rates and Frequencies in Ca professional landscape management pla or area. A suggested application rate is a applied every 4 to 8 weeks during the loc <b>Application Rates:</b> To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of pro- To feed at the rate of 0.12 kg Nitrogen (I	nada: Follow your n for your local cours 1 to 2.8 Kg for 100 m <sup>2</sup> cal growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg
ebanon brand fertilizers are formulated for profession etermined by the professional applicator based on lo Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre.	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlim Wash hands and exposed skin thoroug Do not eat, drink or smoke when using t Wear protective gloves/protective clothi	ate. Arning y 1y after handling. his product. ng/eye protection/face protection.	Application Rates and Frequencies in Ca professional landscape management pla or area. A suggested application rate is a applied every 4 to 8 weeks during the loc <b>Application Rates:</b> To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> 0). Equivalent to 2.0 kg of pr To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> 0). Equivalent to 1.0 kg of pr	nada: Follow your n for your local cours 1 to 2.8 Kg for 100 m <sup>2</sup> cal growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> .
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre. SUGESTED SPREADER SETTINGS	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlin Wash hands and exposed skin thorougt Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek med	ate. Arning y 1y after handling. his product. ng/eye protection/face protection.	Application Rates and Frequencies in Ca professional landscape management pla or area. A suggested application rate is t applied every 4 to 8 weeks during the loc <b>Application Rates:</b> To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> 0). Equivalent to 2.0 kg of pr To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> 0). Equivalent to 1.0 kg of pr <b>SUGGESTED SPREADER</b>	nada: Follow your n for your local cours 1 to 2.8 Kg for 100 m <sup>2</sup> cal growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . SETTINGS
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K <sub>2</sub> 0)/1,000 sq. ft. 0.5# (K <sub>2</sub> 0)/1,000 sq. ft.	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlin Wash hands and exposed skin thorougi Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek met Rinse mouth.	ate. arning y s ny after handling. his product. ng/eye protection/face protection. dical attention if symptoms persist.	Application Rates and Frequencies in Ca professional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc <b>Application Rates:</b> To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of pr To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of pr <b>SUGGESTED SPREADER</b> <b>Spreader</b> 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> C	nada: Follow your n for your local cours 1 to 2.8 Kg for 100 m <sup>2</sup> cal growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . SETTINGS D.24 kg K <sub>2</sub> 0 /100 m <sup>2</sup>
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K,0)/1,000 sq. ft. 0.5 # (K,0)/1,000 sq. ft. Lebanon Turf 2 1½	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlin Wash hands and exposed skin thorougf Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and was	ate. arning g hy after handling. his product. ng/eye protection/face protection. dical attention if symptoms persist. h before reuse.	Application Rates and Frequencies in Calprofessional landscape management platering or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc     Application Rates:     To feed at the rate of 0.24 kg Nitrogen (IPotash (K <sub>2</sub> O). Equivalent to 2.0 kg of pr     To feed at the rate of 0.12 kg Nitrogen (IPotash (K <sub>2</sub> O). Equivalent to 1.0 kg of pr     Spreader   0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> C     Lebanon Turf   2	nada: Follow your n for your local cours 1 to 2.8 Kg for 100 m <sup>2</sup> cal growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . SETTINGS J.24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> 1½
ebanon brand fertilizers are formulated for profession letermined by the professional applicator based on local professional applicator based on local per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K_0) per 1,000 sq. ft., apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K_0)/1,000 sq. ft. 0.5# (K_0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlin Wash hands and exposed skin thorough Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and was IF SWALLOWED: Call a POISON CENTE	ate. arning g hy after handling. his product. ng/eye protection/face protection. dical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell.	Application Rates and Frequencies in Ca professional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc <b>Application Rates:</b> To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of pr To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of pr <b>SUGGESTED SPREADER</b> <b>Spreader</b> 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> C Lebanon Turf 2 Andersons AccuPro F	nada: Follow your n for your local cours t to 2.8 Kg for 100 m <sup>2</sup> cal growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . SETTINGS 0.24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> 1 <sup>1</sup> / <sub>2</sub> D
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K,0)/1,000 sq. ft. 0.5# (K,0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlin, Wash hands and exposed skin thorougf Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek mer Rinse mouth. Take off contaminated clothing and was IF SWALLOWED: Call a POISON CENTE IF ON SKIN: Wash with plenty of soap an	ate. arning y y y y after handling. this product. ng/eye protection/face protection. dical attention if symptoms persist. the before reuse. R or doctor/physician if you feel unwell. nd water.	Application Rates and Frequencies in Ca professional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc Application Rates: To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of pr To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of pr SUGGESTED SPREADER Spreader 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> C Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11	nada: Follow your n for your local cours 1 to 2.8 Kg for 100 m <sup>2</sup> cal growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . SETTINGS 0.24 kg K,0 /100 m <sup>2</sup> 1½ D 9
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft. or 105. lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft. or 105. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K <sub>2</sub> 0)/1,000 sq. ft. 0.5 # (K <sub>2</sub> 0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9 Gandy 22 18	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlin Wash hands and exposed skin thorough Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and was IF SWALLOWED: Call a POISON CENTE	ate. arning y y y y after handling. this product. ng/eye protection/face protection. dical attention if symptoms persist. the before reuse. R or doctor/physician if you feel unwell. nd water.	Application Rates and Frequencies in Caprofessional landscape management pla or area. A suggested application rate is applied every 4 to 8 weeks during the loc     Application Rates:     To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of prot	nada: Follow your n for your local cours 1 to 2.8 Kg for 100 m <sup>2</sup> cal growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . SETTINGS 0.24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> 1½ D 9 18
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 2.1 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K <sub>2</sub> 0)/1,000 sq. ft. 0.5 # (K <sub>2</sub> 0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9 Gandy 22 18 Lely 3 II 2 II	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlin, Wash hands and exposed skin thorougf Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek mer Rinse mouth. Take off contaminated clothing and was IF SWALLOWED: Call a POISON CENTE IF ON SKIN: Wash with plenty of soap an	ate. arning y y y after handling. his product. ng/eye protection/face protection. dical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. nd water. doctor/physician if you feel unwell.	Application Rates and Frequencies in Caprofessional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc     Application Rates:     To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of prime To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of prime SUGGESTED SPREADER Spreader 0.48 kg K <sub>2</sub> O /100 m² C Lebanon Turf 2     Andersons AccuPro   F     Earthway Rotary   11     Gandy   22     Lely   3 II	nada: Follow your n for your local cours 1 to 2.8 Kg for 100 m <sup>2</sup> cal growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . SETTINGS 0.24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> 1½ D 9 18 2 II
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) per 1,000 sq. ft. apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K,0)/1,000 sq. ft. 0.5# (K,0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9 Gandy 22 18 Lely 3 ll 2 ll Lesco (letter/numeric)E/15 D/14	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlim Wash hands and exposed skin thorougi Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and was IF SWALLOWED: Call a POISON CENTE IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER or	ate. arning y y y y y y y y y y y y y	Application Rates and Frequencies in Ca professional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc <b>Application Rates:</b> To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of pr To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of pr <b>SUGGESTED SPREADER</b> <b>Spreader</b> 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> O Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11 Gandy 22 Lely 3 II Lesco (letter/numeric)E/15	nada: Follow your n for your local cours t to 2.8 Kg for 100 m <sup>2</sup> cal growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . SETTINGS 0.24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> 1½ D 9 18 2 II D/14
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) per 1,000 sq. ft. apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K,0)/1,000 sq. ft. 0.5# (K,0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9 Gandy 22 18 Lely 3 II 2 II Lesco (letter/numeric)E/15 D/14 PennMulch HV0 G D	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlin Wash hands and exposed skin thoroug! Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and was IF SWALLOWED: Call a POISON CENTE IF ON SKIN: Wash with plenty of soap au IF INHALED: Call » POISON CENTER or If skin irritation occurs: Get medical adv If eye irritation persists: Get medical adv	ate. arning y y y y y y y y y y y y y	Application Rates and Frequencies in Ca professional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc Application Rates: To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> 0). Equivalent to 2.0 kg of pr To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> 0). Equivalent to 1.0 kg of pr SUGGESTED SPREADER Spreader 0.48 kg K <sub>2</sub> 0 /100 m <sup>2</sup> (I Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11 Gandy 22 Lely 3 II Lesco (letter/numeric)E/15 PennMulch HV0 G	nada: Follow your n for your local cours: 1 to 2.8 Kg for 100 m <sup>2</sup> cal growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . SETTINGS J.24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> J <sup>1</sup> / <sub>2</sub> D 9 18 2 II D/14 D
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 105 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K,0)/1,000 sq. ft. 0.5 # (K,0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9 Gandy 22 18 Lely 3 II 2 II Lesco (letter/numeric)E/15 D/14 PennMulch HV0 G D ProScape SS G D	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlin Wash hands and exposed skin thorougt Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and was IF SWALLOWED: Call a POISON CENTE IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER or If skin irritation occurs: Get medical add If eye irritation persists: Get medical add Store locked up.	ate. arning y y y y y y y y y y y y y	Application Rates and Frequencies in Caprofessional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc     Application Rates:     To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of pr	nada: Follow your n for your local cours: 1 to 2.8 Kg for 100 m <sup>2</sup> cal growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . SETTINGS 0.24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> 1½ 0 9 18 2 II 0/14 0 0 0
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft. apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K <sub>2</sub> 0)/1,000 sq. ft. 0.5 # (K <sub>2</sub> 0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9 Gandy 22 18 Lely 3 II 2 II Lesco (letter/numeric)E/15 D/14 PennMulch HV0 G D ProScape SS G D Spyker 2 1½	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlin Wash hands and exposed skin thorougi Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek mer Rinse mouth. Take off contaminated clothing and was IF SWALLOWED: Call a POISON CENTE IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER or If skin irritation occurs: Get medical adu Store locked up. Store in a well-ventilated place.	ate. arning y y y y y y y y y y y y y	Application Rates and Frequencies in Caprofessional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc     Application Rates:     To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent to 1.0 kg of protosh (K <sub>2</sub> O). Equivalent (K <sub>2</sub> O). Equivalent (K <sub>2</sub> O). Equivalent (K <sub>2</sub> O). Equival	nada: Follow your n for your local cours: 1 to 2.8 Kg for 100 m <sup>2</sup> cal growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . SETTINGS 0.24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> 1½ D 9 18 2 II D/14 D D 1½
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K_0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K_0) per 1,000 sq. ft., apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K_0)/1,000 sq. ft. 0.5# (K_0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9 Gandy 22 18 Lely 3 II 2 1I Lesco (letter/numeric)E/15 D/14 PennMulch HV0 G D ProScape SS G D Spyker 2 1½	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlin, Wash hands and exposed skin thorougf Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek mer Rinse mouth. Take off contaminated clothing and was IF SWALLOWED: Call a POISON CENTE IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER or If skin irritation persists: Get medical adv If eye irritation persists: Get medical adv Store locked up. Store in a well-ventilated place. Keep container tightly closed.	ate. arning y after handling. this product. ng/eye protection/face protection. dical attention if symptoms persist. the before reuse. R or doctor/physician if you feel unwell. nd water. doctor/physician if you feel unwell. rice/ attention. wice/attention.	Application Rates and Frequencies in Caprofessional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc     Application Rates:     To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of pr     To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of pr     To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of pr     Spreader   0.48 kg K <sub>2</sub> O /100 m² O     Lebanon Turf   2     Andersons AccuPro   F     Earthway Rotary   11     Gandy   22     Lely   3 II     Lesco (letter/numeric)E/15   PennMulch HVO     ProScape SS   G     Spyker   2     Vicon   17	nada: Follow your n for your local course t to 2.8 Kg for 100 m <sup>2</sup> , cal growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . <b>SETTINGS</b> <b>D.24 kg K<sub>2</sub>0 /100 m<sup>2</sup></b> 1½ D 9 18 2 II D/14 D 1½ 14
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K,0)/1,000 sq. ft. 0.5# (K,0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9 Gandy 22 18 Lely 3 II 2 II Lesco (letter/numeric)E/15 D/14 PennMulch HV0 G D ProScape SS G D Spyker 2 1½ Vicon 17 14 These settings were calibrated and field tested.	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlin, Wash hands and exposed skin thorougf Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek mer Rinse mouth. Take off contaminated clothing and was IF SWALLOWED: Call a POISON CENTE IF ON SKIN: Wash with plenty of soap an IF SWALLOWED: Call a POISON CENTER or If skin irritation occurs: Get medical adv If eye irritation persists: Get medical adv Store in a well-ventilated place. Keep container tightly closed. Dispose of contents/container property	ate. arning y s y y y after handling. this product. ng/eye protection/face protection. dical attention if symptoms persist. the before reuse. R or doctor/physician if you feel unwell. nd water. doctor/physician if you feel unwell. rice/ attention. wice/attention.	Application Rates and Frequencies in Caprofessional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc     Application Rates:     To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> 0). Equivalent to 2.0 kg of pr     To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> 0). Equivalent to 1.0 kg of pr     To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> 0). Equivalent to 1.0 kg of pr     Spreader   0.48 kg K <sub>2</sub> 0 /100 m <sup>2</sup> O     Lebanon Turf   2     Andersons AccuPro   F     Earthway Rotary   11     Gandy   22     Lely   3 II     Lesco (letter/numeric)E/15   PennMulch HVO     ProScape SS   6     Spyker   2     Vicon   17     These settings were calibrated and	nada: Follow your n for your local course t to 2.8 Kg for 100 m <sup>2</sup> cal growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . <b>SETTINGS</b> <b>J.24 kg K<sub>2</sub>0 /100 m<sup>2</sup></b> <b>J</b> <sup>1</sup> / <sub>2</sub> <b>D</b> <b>9</b> 18 2 II D/14 D <b>1</b> <i>1</i> / <sub>2</sub> 14 d field tested.
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K,0)/1,000 sq. ft. 0.5# (K,0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9 Gandy 22 18 Lely 3 II 2 II Lesco (letter/numeric)E/15 D/14 PennMulch HV0 G D ProScape SS G D Spyker 2 1½ Vicon 17 14 These settings were calibrated and field tested. However, age and condition of spreader, speed of	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlin Wash hands and exposed skin thoroug Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and was IF SWALLOWED: Call a POISON CENTE IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER or If skin irritation occurs: Get medical adv Store locked up. Store in a well-ventilated place. Keep container tightly closed. Dispose of contents/container properly. Keep out of reach of children	ate. arning y y y y y y y y y y y y y	Application Rates and Frequencies in Caprofessional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc     Application Rates:     To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> 0). Equivalent to 2.0 kg of pr     To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> 0). Equivalent to 1.0 kg of pr     SUGGESTED SPREADER     Spreader   0.48 kg K <sub>2</sub> 0 /100 m² (I Lebanon Turf     2   Andersons AccuPro     Farthway Rotary   11     Gandy   22     Lely   3 II     Lesco (letter/numeric)E/15     PennMulch HVO   6     ProScape SS   6     Spyker   2     Vicon   17     These settings were calibrated and However, age and condition of spr	nada: Follow your n for your local cours: 1 to 2.8 Kg for 100 m <sup>2</sup> cal growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . SETTINGS 0.24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> 1½ 0 9 18 2 II 0/14 0 0 1½ 14 d field tested. eader, speed of
ebanon brand fertilizers are formulated for profession letermined by the professional applicator based on local Particular and the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K,0)/1,000 sq. ft. 0.5 # (K,0)/1,000 sq. ft. Lebanon Turf 2 $1\frac{1}{2}$ Andersons AccuPro F D Earthway Rotary 11 9 Gandy 22 18 Lely 3 II 2 II Lesco (letter/numeric)E/15 D/14 PennMulch HV0 G D ProScape SS G D Spyker 2 $1\frac{1}{2}$ Vicon 17 14 These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlin Wash hands and exposed skin thoroug! Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and was IF SWALLOWED: Call a POISON CENTER IF ON SKIN: Wash with plenty of soap an IF INHALED: Call a POISON CENTER or If skin irritation occurs: Get medical adv Store locked up. Store in a well-ventilated place. Keep container tightly closed. Dispose of contents/container properly Keep out of reach of children	ate. arning y s y y y after handling. this product. ng/eye protection/face protection. dical attention if symptoms persist. the before reuse. R or doctor/physician if you feel unwell. nd water. doctor/physician if you feel unwell. rice/ attention. wice/attention.	Application Rates and Frequencies in Caprofessional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc     Application Rates:     To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> 0). Equivalent to 2.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of protosh (K <sub>2</sub> 0). Equivalent to 1.0 kg of pr	nada: Follow your n for your local cours: 1 to 2.8 Kg for 100 m <sup>2</sup> cal growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . SETTINGS 0.24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> 1½ 0 9 18 2 II D/14 0 0 1½ 14 d field tested. eader, speed of may require slightly
ebanon brand fertilizers are formulated for profession letermined by the professional applicator based on location by the profession by the profess	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlin Wash hands and exposed skin thorougt Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and was IF SWALLOWED: Call a POISON CENTE IF ON SKIN: Wash with plenty of soap an IF INHALED: Call a POISON CENTER or If skin irritation occurs: Get medical add Store locked up. Store in a well-ventilated place. Keep container tightly closed. Dispose of contents/container properly. Keep out of reach of children Precautionary Statements for disposa state and local regulations.	ate. arning y any after handling. his product. ng/eye protection/face protection. dical attention if symptoms persist. the before reuse. R or doctor/physician if you feel unwell. nd water. doctor/physician if you feel unwell. ice/ attention. vice/attention. vice/attention. as required by law for waste pesticides. I - Dispose in accordance with all federal,	Application Rates and Frequencies in Caprofessional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc     Application Rates:     To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of pr     To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of pr     SUGGESTED SPREADER     Spreader   0.48 kg K <sub>2</sub> O /100 m² O     Lebanon Turf   2     Andersons AccuPro   F     Earthway Rotary   11     Gandy   22     Lely   3 II     Lesco (letter/numeric)E/15     PensMulch HVO   G     ProScape SS   G     Spyker   2     Vicon   17     These settings were calibrated and However, age and condition of spreoperation and evenness of terrain different settings for desired cover	nada: Follow your n for your local course 1 to 2.8 Kg for 100 m <sup>2</sup> , al growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . <b>SETTINGS</b> <b>0.24 kg K_0 /100 m<sup>2</sup></b> <b>1%</b> <b>1%</b> <b>18</b> <b>2 II</b> <b>D</b> /14 <b>D</b> <b>1%</b> <b>14</b> <b>d field tested.</b> eader, speed of may require slightly
ebanon brand fertilizers are formulated for profession etermined by the professional applicator based on loging etermined by the profession etermined by the p	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlin Wash hands and exposed skin thorougt Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek meet Rinse mouth. Take off contaminated clothing and was IF SWALLOWED: Call a POISON CENTE IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER or If skin irritation occurs: Get medical add Store locked up. Store in a well-ventilated place. Keep container tightly closed. Dispose of contents/container properly. Keep out of reach of children Precautionary Statements for disposa state and local regulations. Fertilizer may cause irritation of eye	ate. arning y g hy after handling. his product. ng/eye protection/face protection. dical attention if symptoms persist. the before reuse. R or doctor/physician if you feel unwell. nd water. doctor/physician if you feel unwell. ice/ attention. vice/attention. as required by law for waste pesticides. I - Dispose in accordance with all federal, es, nose, throat, and skin. In case of com	Application Rates and Frequencies in Caprofessional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc     Application Rates:     To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of pr     To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of pr     SUGGESTED SPREADER     Spreader   0.48 kg K <sub>2</sub> O /100 m² O     Lebanon Turf   2     Andersons AccuPro   F     Earthway Rotary   11     Gandy   22     Lely   3 II     Lesco (letter/numeric)E/15     PensMulch HVO   G     ProScape SS   G     Spyker   2     Vicon   17     These settings were calibrated and However, age and condition of spreoperation and evenness of terrain different settings for desired cover	nada: Follow your n for your local course 1 to 2.8 Kg for 100 m <sup>2</sup> , al growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . <b>SETTINGS</b> <b>0.24 kg K_0 /100 m<sup>2</sup></b> <b>1%</b> <b>1%</b> <b>18</b> <b>2 II</b> <b>D</b> /14 <b>D</b> <b>1%</b> <b>14</b> <b>d field tested.</b> eader, speed of may require slightly
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) per 1,000 sq. ft. apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K,0)/1,000 sq. ft. 0.5# (K,0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9 Gandy 22 18 Lely 3 II 2 II Lesco (letter/numeric)E/15 D/14 PennMulch HV0 G D ProScape SS G D Spyker 2 1½ Vicon 17 14 These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlin Wash hands and exposed skin thorougt Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and was IF SWALLOWED: Call a POISON CENTE IF ON SKIN: Wash with plenty of soap an IF INHALED: Call a POISON CENTER or If skin irritation occurs: Get medical add Store locked up. Store in a well-ventilated place. Keep container tightly closed. Dispose of contents/container properly. Keep out of reach of children Precautionary Statements for disposa state and local regulations.	ate. arning y g hy after handling. his product. ng/eye protection/face protection. dical attention if symptoms persist. the before reuse. R or doctor/physician if you feel unwell. nd water. doctor/physician if you feel unwell. ice/ attention. vice/attention. as required by law for waste pesticides. I - Dispose in accordance with all federal, es, nose, throat, and skin. In case of com	Application Rates and Frequencies in Caprofessional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc     Application Rates:     To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of pr     To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of pr     SUGGESTED SPREADER     Spreader   0.48 kg K <sub>2</sub> O /100 m² O     Lebanon Turf   2     Andersons AccuPro   F     Earthway Rotary   11     Gandy   22     Lely   3 II     Lesco (letter/numeric)E/15     PensMulch HVO   G     ProScape SS   G     Spyker   2     Vicon   17     These settings were calibrated and However, age and condition of spreoperation and evenness of terrain different settings for desired cover	nada: Follow your n for your local cours 1 to 2.8 Kg for 100 m <sup>2</sup> cal growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . SETTINGS 0.24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> 1½ 0 9 18 2 II 0/14 0 0 1½ 14 d field tested. eader, speed of may require slightly
ebanon brand fertilizers are formulated for profession letermined by the professional applicator based on local etermined by the professional applicator based on local application Rates:     To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre.     SUGGESTED SPREADER SETTINGS     Spreader   1½     Andersons AccuPro   F     D   D     Earthway Rotary   11   9     Gandy   22   18     Lely   3 ll   2 ll     Lesco (letter/numeric)E/15   D/14     PensMulch HVO   G   D     ProScape SS   G   D     Spyker   2   1½     Vicon   17   14     These settings were calibrated and field tested.   However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorougf Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and wass IF SWALLOWED: Call a POISON CENTER or If skin irritation occurs: Get medical adv Store locked up. Store in a well-ventilated place. Keep container tightly closed. Dispose of contents/container properly Keep out of reach of children Precautionary Statements for disposa state and local regulations.	ate. arning y y after handling. this product. ng/eye protection/face protection. dical attention if symptoms persist. th before reuse. R or doctor/physician if you feel unwell. nd water. doctor/physician if you feel unwell. ice/ attention. vice/attention. as required by law for waste pesticides. I - Dispose in accordance with all federal, es, nose, throat, and skin. In case of con- tter; for eyes get medical attention.	Application Rates and Frequencies in Ca professional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc Application Rates: To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of pr To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of pr To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of pr SUGGESTED SPREADER Spreader 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> C Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11 Gandy 22 Lely 3 II Lesco (letter/numeric)E/15 PennMulch HVO G ProScape SS G Spyker 2 Vicon 17 These settings were calibrated and However, age and condition of spro operation and evenness of terrain different settings for desired cove tact with	nada: Follow your n for your local course 1 to 2.8 Kg for 100 m <sup>2</sup> , al growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . <b>SETTINGS</b> <b>0.24 kg K_0 /100 m<sup>2</sup></b> <b>1%</b> <b>1%</b> <b>18</b> <b>2 II</b> <b>D</b> /14 <b>D</b> <b>1%</b> <b>14</b> <b>d field tested.</b> eader, speed of may require slightly
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) per 1,000 sq. ft. apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K,0)/1,000 sq. ft. 0.5# (K,0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9 Gandy 22 18 Lely 3 II 2 II Lesco (letter/numeric)E/15 D/14 PennMulch HV0 G D ProScape SS G D Spyker 2 1½ Vicon 17 14 These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handlin Wash hands and exposed skin thorougi Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and was IF SWALLOWED: Call a POISON CENTER IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER or If skin irritation occurs: Get medical adv Store locked up. Store in a well-ventilated place. Keep container tightly closed. Dispose of contents/container properly. Keep out of reach of children Precautionary Statements for disposa state and local regulations. Fertilizer may cause irritation of eye skin or eyes, flush with plenty of wa	ate. arning y y after handling. his product. ng/eye protection/face protection. dical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. nd water. doctor/physician if you feel unwell. rice/ attention. vice/attention. as required by law for waste pesticides. I - Dispose in accordance with all federal, as, nose, throat, and skin. In case of com ter; for eyes get medical attention. lations of a qualified individual or institu	Application Rates and Frequencies in Ca professional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc Application Rates: To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> 0). Equivalent to 2.0 kg of pr To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> 0). Equivalent to 1.0 kg of pr To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> 0). Equivalent to 1.0 kg of pr SUGGESTED SPREADER Spreader 0.48 kg K <sub>2</sub> 0 /100 m <sup>2</sup> (I Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11 Gandy 22 Lely 3 ll Lesco (letter/numeric)E/15 PennMulch HV0 G ProScape SS G Spyker 2 Vicon 17 These settings were calibrated and However, age and condition of spro operation and evenness of terrain different settings for desired cove tact with	nada: Follow your n for your local course 1 to 2.8 Kg for 100 m <sup>2</sup> , al growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . <b>SETTINGS</b> <b>0.24 kg K_0 /100 m<sup>2</sup></b> <b>1%</b> <b>1%</b> <b>18</b> <b>2</b> II <b>D/14</b> <b>D</b> <b>1%</b> <b>14</b> <b>d field tested.</b> eader, speed of may require slightly
Application Rates:     To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) oer 1,000 sq. ft. apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K,0) per 1,000 sq. ft. apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre.     SUGGESTED SPREADER SETTINGS     Spreader   1½     Andersons AccuPro   D     Earthway Rotary   11   9     Gandy   22   18     Lely   3 ll   2 ll     Lesco (letter/numeric)E/15   D/14     PronMulch HVO   G   D     ProScape SS   G   D     Spyker   2   1½     These settings were calibrated and field tested.   However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and was IF SWALLOWED: Call a POISON CENTE IF ON SKIN: Wash with plenty of soap an IF INHALED: Call a POISON CENTER or If skin irritation occurs: Get medical adv Store locked up. Store in a well-ventilated place. Keep container tightly closed. Dispose of contents/container properly. Keep out of reach of children Precautionary Statements for disposa state and local regulations. Fertilizer may cause irritation of eye skin or eyes, flush with plenty of wa	ate. arning y y after handling. his product. ng/eye protection/face protection. dical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. nd water. doctor/physician if you feel unwell. rice/ attention. vice/attention. is required by law for waste pesticides. I - Dispose in accordance with all federal, es, nose, throat, and skin. In case of com iter; for eyes get medical attention. lations of a qualified individual or institut d crop advisor, agronomist, university c	Application Rates and Frequencies in Ca professional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc Application Rates: To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of pr To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of pr To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of pr SUGGESTED SPREADER Spreader 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> C Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11 Gandy 22 Lely 3 II Lesco (letter/numeric)E/15 PennMulch HVO G ProScape SS G Spyker 2 Vicon 17 These settings were calibrated and However, age and condition of spro operation and evenness of terrain different settings for desired cove tact with	nada: Follow your n for your local course 1 to 2.8 Kg for 100 m <sup>2</sup> , al growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . <b>SETTINGS</b> <b>0.24 kg K_0 /100 m<sup>2</sup></b> <b>1%</b> <b>1%</b> <b>18</b> <b>2</b> II <b>D/14</b> <b>D</b> <b>1%</b> <b>14</b> <b>d field tested.</b> eader, speed of may require slightly
Application Rates:     To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) ore 1,000 sq. ft. apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K,0) per 1,000 sq. ft. apply this product at 2.1 bs. per 1,000 sq. ft. or 120 lbs. per acre.     SUGGESTED SPREADER SETTINGS     Spreader   1½     Andersons AccuPro   D     Earthway Rotary   11   9     Bandy   22   18     Lely   3 ll   2 ll     Lesco (letter/numeric)E/15   D/14     PronMulch HVO   G   D     ProScape SS   G   D     Spyker   2   1½     These settings were calibrated and field tested.   However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.	nal use on turf, including golf c cal soils, regulations, and clim Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using t Wear protective gloves/protective clothi If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and was IF SWALLOWED: Call a POISON CENTE IF ON SKIN: Wash with plenty of soap an IF INHALED: Call a POISON CENTER or If skin irritation occurs: Get medical adv Store locked up. Store in a well-ventilated place. Keep container tightly closed. Dispose of contents/container properly. Keep out of reach of children Precautionary Statements for disposa state and local regulations. Fertilizer may cause irritation of eye skin or eyes, flush with plenty of wa	ate. arning y y after handling. his product. ng/eye protection/face protection. dical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. nd water. doctor/physician if you feel unwell. rice/ attention. vice/attention. as required by law for waste pesticides. I - Dispose in accordance with all federal, as, nose, throat, and skin. In case of com ter; for eyes get medical attention. lations of a qualified individual or institu	Application Rates and Frequencies in Ca professional landscape management pla or area. A suggested application rate is 1 applied every 4 to 8 weeks during the loc Application Rates: To feed at the rate of 0.24 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of pr To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of pr To feed at the rate of 0.12 kg Nitrogen (I Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of pr SUGGESTED SPREADER Spreader 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> C Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11 Gandy 22 Lely 3 II Lesco (letter/numeric)E/15 PennMulch HVO G ProScape SS G Spyker 2 Vicon 17 These settings were calibrated and However, age and condition of spro operation and evenness of terrain different settings for desired cove tact with	nada: Follow your n for your local cours 1 to 2.8 Kg for 100 m <sup>2</sup> cal growing season. N) or 0.48 kg oduct per 100 m <sup>2</sup> . N) or 0.24 kg oduct per 100 m <sup>2</sup> . SETTINGS 0.24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> 1½ 0 9 18 2 II 0/14 0 0 1½ 14 d field tested. eader, speed of may require slight!

#### **GUARANTEED MINIMUM ANALYSIS**

88685 35736

F699

21-35736

UUANANTEED ANALIS	010	uuanan	ILLD MIMIMUM ANALISI	
Total Nitrogen (N)	12%	Total Nitrogen (N)		12%
0.9% Ammoniacal Nitrogen		0.9% Ammoniacal Nitrogen		
2.4% Water Insoluble Nitrogen*		2.4% Water Insoluble Nitrog	jen*	
4.5% Urea Nitrogen		4.5% Urea Nitrogen		
4.2% Other Water Soluble Nitrogen *		4.2% Other Water Soluble N		
Soluble Potash (K <sub>2</sub> O)		Soluble Potash (K <sub>2</sub> O)		24%
Magnesium (Mg)	0.5%	Magnesium (Mg)		0.5%
0.5% Water Soluble Magnesium (Mg)		0.5% Water Soluble Magnes	sium (Mg)	
Sulfur (S)		Sulfur (S)		9.5%
9.5% Combined Sulfur (S)		9.5% Combined Sulfur (S)		
Iron (Fe)	1.6%	Iron (Fe) (actual)		1.6%
0.2% Water Soluble Iron (Fe)		0.2% Water Soluble Iron (Fe	e)	
Manganese (Mn)	0.5%	Manganese (Mn)	,	0.5%
0.5% Water Soluble Manganese (Mn)		0.5% Water Soluble Mangar	nese (Mn)	
Derived From: Ammonium Sulfate, Methylene Ureas, Ur of Potash Magnesia, Kelp Meal, Ferric Oxide, Ferrous S		Chlorine (Cl) (maximum) *6.6% Slowly Available Nitrog	gen from Methylene Ureas.	2.0%
*6.6% Slowly Available Nitrogen from Methylene Ureas			ulfate, Methylene Ureas, Urea, Sulfate of F	
Chlorine (CI) not more than		of Potash Magnesia, Kelp Me	al, Ferric Oxide, Ferrous Sulfate, Mangan	ese Sulfate.
ALSO CONTAINS NON PLANT FOOD INGREDIE	ENTS		PLANT FOOD INGREDIENTS	
SOIL/PLANT AMENDING INGREDIENTS		SOIL/PLANT AMENDING	G INGREDIENTS	
Active Ingredients:		Active Ingredients:		
Humic Acids (derived from Leonardite) 1.09			rom Leonardite) 1.0%	CANAD
Inert Ingredients (as soil amending)99.0% 99.0% Fertilizer Ingredients	<sup>/0</sup> U.J.A	Inert Ingredients (as so 99.0% Fertilizer In	oil amending)99.0%	CANAD
	<del>b© MD products contain humic</del>	1	•	
otice: This product contains the micronutrient iron. Iron may st romptly after applications by sweeping or blowing. Do not wash ebanon brand fertilizers are formulated for professio	ı off with water. nal use on turf, including golf cı	be applied on dry or water dampened o ourses, sports fields, and landso	concrete and should be removed from these ar	
otice: This product contains the micronutrient iron. Iron may st romptly after applications by sweeping or blowing. Do not wash	ı off with water. nal use on turf, including golf cı	be applied on dry or water dampened o ourses, sports fields, and landso	concrete and should be removed from these ar	requencies are
otice: This product contains the micronutrient iron. Iron may st romptly after applications by sweeping or blowing. Do not wash ebanon brand fertilizers are formulated for professio	n off with water. nal use on turf, including golf co ocal soils, regulations, and clima Wa	be applied on dry or water dampened o ourses, sports fields, and landso ate.	concrete and should be removed from these ar capes. Specific application rates and f Application Rates and Frequencies in Cana	requencies are
otice: This product contains the micronutrient iron. Iron may st romptly after applications by sweeping or blowing. Do not wash ebanon brand fertilizers are formulated for professio	n off with water. nal use on turf, including golf co ocal soils, regulations, and clima Hazard Statements Wa	be applied on dry or water dampened o ourses, sports fields, and landso	concrete and should be removed from these ar capes. Specific application rates and f Application Rates and Frequencies in Cana professional landscape management plan	<b>requencies are</b> ada: Follow your for your local cours
otice: This product contains the micronutrient iron. Iron may st romptly after applications by sweeping or blowing. Do not wash ebanon brand fertilizers are formulated for professio	n off with water. nal use on turf, including golf co ocal soils, regulations, and clima Wa	be applied on dry or water dampened o ourses, sports fields, and landso ate.	concrete and should be removed from these ar capes. Specific application rates and f Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t	<b>requencies are</b> ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup>
otice: This product contains the micronutrient iron. Iron may st romptly after applications by sweeping or blowing. Do not wash ebanon brand fertilizers are formulated for professio etermined by the professional applicator based on lo	n off with water. nal use on turf, including golf co cal soils, regulations, and clima Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation.	be applied on dry or water dampened o ourses, sports fields, and landso ate.	concrete and should be removed from these ar capes. Specific application rates and f Application Rates and Frequencies in Cana professional landscape management plan	<b>requencies are</b> ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup>
otice: This product contains the micronutrient iron. Iron may st romptly after applications by sweeping or blowing. Do not wash ebanon brand fertilizers are formulated for professio etermined by the professional applicator based on lo Application Rates:	n off with water. nal use on turf, including golf co ocal soils, regulations, and clima Hazard Statements Harmful if swallowed. Causes skin irritation. Causes seye irritation. May be harmful if inhaled.	be applied on dry or water dampened o ourses, sports fields, and landso ate.	concrete and should be removed from these ar capes. Specific application rates and f Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local	<b>requencies are</b> ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup>
otice: This product contains the micronutrient iron. Iron may st romptly after applications by sweeping or blowing. Do not wash ebanon brand fertilizers are formulated for profession etermined by the professional applicator based on lo Application Rates: Fo feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0)	n off with water. nal use on turf, including golf co cal soils, regulations, and clima Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation.	be applied on dry or water dampened of ourses, sports fields, and landso ate. Irning	concrete and should be removed from these ar capes. Specific application rates and f Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local Application Rates:	<b>requencies are</b> ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> I growing season.
otice: This product contains the micronutrient iron. Iron may st romptly after applications by sweeping or blowing. Do not wash ebanon brand fertilizers are formulated for profession etermined by the professional applicator based on lo Application Rates: Fo feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) ber 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or	n off with water. nal use on turf, including golf co incal soils, regulations, and clima Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling	be applied on dry or water dampened of ourses, sports fields, and landso ate. Irning	concrete and should be removed from these ar capes. Specific application rates and f Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> l growing season. o or 0.48 kg
otice: This product contains the micronutrient iron. Iron may st romptly after applications by sweeping or blowing. Do not wash ebanon brand fertilizers are formulated for profession etermined by the professional applicator based on lo Application Rates: Fo feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or	n off with water. nal use on turf, including golf co total soils, regulations, and clima Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough	be applied on dry or water dampened o ourses, sports fields, and landso ate. Irning I after handling.	concrete and should be removed from these are capes. Specific application rates and for Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local Application Rates: To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> I growing season. or 0.48 kg Juct per 100 m <sup>2</sup> .
otice: This product contains the micronutrient iron. Iron may st romptly after applications by sweeping or blowing. Do not wash ebanon brand fertilizers are formulated for profession etermined by the professional applicator based on lo Application Rates: Fo feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 2.1	n off with water. nal use on turf, including golf co total soils, regulations, and clima Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th	be applied on dry or water dampened o ourses, sports fields, and landso ate. Irning J y after handling. his product.	concrete and should be removed from these are capes. Specific application rates and for Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local Application Rates: To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N)	requencies are da: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> l growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg
otice: This product contains the micronutrient iron. Iron may st romptly after applications by sweeping or blowing. Do not wash ebanon brand fertilizers are formulated for profession etermined by the professional applicator based on lo profeed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 83 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 2.1 bs. per 1,000 sq. ft. or 120 lbs. per acre.	n off with water. nal use on turf, including golf co total soils, regulations, and clima Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th Wear protective gloves/protective clothin	be applied on dry or water dampened of ourses, sports fields, and landsd ate. Irming Iv after handling. his product. ng/eye protection/face protection.	concrete and should be removed from these are capes. Specific application rates and for Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local <b>Application Rates:</b> To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of proc	requencies are da: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> l growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg duct per 100 m <sup>2</sup> .
otice: This product contains the micronutrient iron. Iron may st romptly after applications by sweeping or blowing. Do not wash ebanon brand fertilizers are formulated for profession etermined by the professional applicator based on lo Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 1.5 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 2.1 bs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS	n off with water. nal use on turf, including golf con- total soils, regulations, and clima Hazard Statements Harmful if swallowed. Causes skin irritation. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th Wear protective gloves/protective clothir If in eyes, rinse with water and seek med	be applied on dry or water dampened of ourses, sports fields, and landsd ate. Irming Iv after handling. his product. ng/eye protection/face protection.	concrete and should be removed from these are capes. Specific application rates and for Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local Application Rates: To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of proc SUGGESTED SPREADER S	requencies are da: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> l growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg duct per 100 m <sup>2</sup> . ETTINGS
otice: This product contains the micronutrient iron. Iron may st romptly after applications by sweeping or blowing. Do not wash ebanon brand fertilizers are formulated for profession etermined by the professional applicator based on lo Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS 1# (K <sub>2</sub> 0)/1,000 sq. ft. 0.5# (K <sub>2</sub> 0)/1,000 sq. ft.	n off with water. nal use on turf, including golf con- total soils, regulations, and clima Hazard Statements Harmful if swallowed. Causes skin irritation. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th Wear protective gloves/protective clothir If in eyes, rinse with water and seek med Rinse mouth.	be applied on dry or water dampened of ourses, sports fields, and landsd ate. Irning Iv after handling. his product. ng/eye protection/face protection. lical attention if symptoms persist.	concrete and should be removed from these are capes. Specific application rates and for Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local Application Rates: To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of proc SUGGESTED SPREADER S Spreader 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> 0.2	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> l growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg duct per 100 m <sup>2</sup> . ETTINGS 24 kg K <sub>2</sub> 0 /100 m <sup>2</sup>
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) Der 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K <sub>2</sub> 0/1,000 sq. ft. 0.5 # (K <sub>2</sub> 0)/1,000 sq. ft. 1# (K <sub>2</sub> 0) 1/2 lbs. per acre. SugGESTED SPREADER SETTINGS Spreader 1# (K <sub>2</sub> 0/1,000 sq. ft. 0.5 # (K <sub>2</sub> 0)/1,000 sq. ft. 1% (K <sub>2</sub> 0)/1,000 sq. ft. 1% (K <sub>2</sub> 0)/1,000 sq. ft.	n off with water. nal use on turf, including golf con- total soils, regulations, and clima Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th Wear protective gloves/protective clothir If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and wast	be applied on dry or water dampened of ourses, sports fields, and landso ate. Irning y after handling. his product. ng/eye protection/face protection. lical attention if symptoms persist. h before reuse.	concrete and should be removed from these are capes. Specific application rates and fr Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local Application Rates: To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of proc SUGGESTED SPREADER S Spreader 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> 0.2	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> I growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg duct per 100 m <sup>2</sup> . ETTINGS 24 kg K <sub>0</sub> 0 /100 m <sup>2</sup> 1 <sup>1</sup> / <sub>2</sub>
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K <sub>2</sub> 0)/1,000 sq. ft. 0.5 f(K <sub>2</sub> 0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D	n off with water. nal use on turf, including golf con- total soils, regulations, and climar Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th Wear protective gloves/protective clothir If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and wast IF SWALLOWED: Call a POISON CENTER	be applied on dry or water dampened of ourses, sports fields, and landso ate. Irning Iv after handling. his product. ng/eye protection/face protection. lical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell.	concrete and should be removed from these are capes. Specific application rates and f Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local Application Rates: To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of proc SUGGESTED SPREADER S Spreader 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> 0.2 Lebanon Turf 2 Andersons AccuPro F	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> l growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg duct per 100 m <sup>2</sup> . ETTINGS 24 kg K <sub>2</sub> 0 /100 m <sup>2</sup>
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) the rate of 0.5 lb. Nitrogen (N) or 1.5 lb. of Potash (K,0) per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K,0)/1,000 sq. ft. 0.5 # (K,0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9 Gandy 22 18	n off with water. nal use on turf, including golf con- total soils, regulations, and clima Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th Wear protective gloves/protective clothir If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and wass IF SWALLOWED: Call a POISON CENTEF IF ON SKIN: Wash with plenty of soap an	be applied on dry or water dampened of ourses, sports fields, and landso ate. Irning Ivafter handling. his product. ng/eye protection/face protection. lical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. nd water.	concrete and should be removed from these are capes. Specific application rates and fr Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local Application Rates: To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of proc SUGGESTED SPREADER S Spreader 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> 0.2	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> l growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg duct per 100 m <sup>2</sup> . ETTINGS 24 kg K0 /100 m <sup>2</sup> D
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) Det 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 1.5 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 2.1 bs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K <sub>2</sub> 0)/1,000 sq. ft. 0.5 (K <sub>2</sub> 0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9 Earthy 22 18 Lely 3 II 2 II	n off with water. nal use on turf, including golf con- total soils, regulations, and clima Hazard Statements Harmful if swallowed. Causes skin irritation. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th Wear protective gloves/protective clothir If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and wash IF SWALLOWED: Call a POISON CENTER IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER or of	be applied on dry or water dampened of ourses, sports fields, and landso ate. Irning Ivafter handling. his product. ng/eye protection/face protection. lical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. d water.	concrete and should be removed from these are capes. Specific application rates and fr Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local Application Rates: To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of proc SUGGESTED SPREADER S Spreader 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> 0.2 Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11 Gandy 22 Lely 3 II	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> l growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg duct per 100 m <sup>2</sup> . ETTINGS 24 kg K <sub>0</sub> 0 /100 m <sup>2</sup> $1\frac{V_2}{9}$ 9
Application Rates:     To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) or 1000 sq. ft. or 120 lbs. per 1,000 sq. ft. or 120 lbs. per 1,000 sq. ft. or 120 lbs. per acre.     SUGGESTED SPREADER SETTINGS     Spreader 1# (K <sub>2</sub> 0)/1,000 sq. ft. 0.5 # (K <sub>2</sub> 0)/1,000 sq. ft. or 120 lbs. per acre.     SUGGESTED SPREADER SETTINGS     Spreader 1# (K <sub>2</sub> 0)/1,000 sq. ft. 0.5 # (K <sub>2</sub> 0)/1,000 sq. ft. 1%2     Andersons AccuPro   F     Deathway Rotary   11   9     Gandy   22   18     Lely   3 II   2 II     Lely   3 II   2 II	n off with water. nal use on turf, including golf co total soils, regulations, and clima Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th Wear protective gloves/protective clothir If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and wash IF SWALLOWED: Call a POISON CENTEF IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTEF or of If skin irritation occurs: Get medical advi	be applied on dry or water dampened of ourses, sports fields, and landsd ate. Irming Jy after handling. his product. ng/eye protection/face protection. lical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. id water. doctor/physician if you feel unwell. ice/ attention.	concrete and should be removed from these are capes. Specific application rates and f Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local Application Rates: To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of proc SUGGESTED SPREADER S Spreader 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> 0.2 Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11 Gandy 22 Lely 3 II Lesso (letter/numeric)E/15	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> l growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg duct per 100 m <sup>2</sup> . ETTINGS 24 kg K,0 /100 m <sup>2</sup> 1 $\frac{1}{2}$ D 9 18 2 II D/14
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft. apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 1.5 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft. apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K <sub>2</sub> 0)/1,000 sq. ft. 0.5 # (K <sub>2</sub> 0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9 Gandy 22 18 Lely 3 II 2 II Lesco (letter/numeric)E/15 D/14 PennMulch HVO G D	n off with water. nal use on turf, including golf co incal soils, regulations, and clima Hazard Statements Harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th Wear protective gloves/protective clothir If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and wast IF SWALLOWED: Call a POISON CENTER IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER or of If skin irritation occurs: Get medical advil If eye irritation persists: Get medical advil	be applied on dry or water dampened of ourses, sports fields, and landsd ate. Irming Jy after handling. his product. ng/eye protection/face protection. lical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. id water. doctor/physician if you feel unwell. ice/ attention.	concrete and should be removed from these are capes. Specific application rates and f Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local <b>Application Rates:</b> To feed at the rate of 0.24 kg Nitrogen (N) Potash ( $K_2$ 0). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N) Potash ( $K_2$ 0). Equivalent to 1.0 kg of proc SUGGESTED SPREADER S Spreader 0.48 kg $K_2$ 0/100 m <sup>2</sup> 0.2 Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11 Gandy 22 Lely 3 II Lesco (letter/numeric)E/15 PennMulch HV0 G	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> l growing season. or 0.48 kg fuct per 100 m <sup>2</sup> . or 0.24 kg fuct per 100 m <sup>2</sup> . ETTINGS 24 kg K,0 /100 m <sup>2</sup> 1½ D 9 18 2 II D/14 D
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K,0) per 1,000 sq. ft. apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K,0) per 1,000 sq. ft., apply this product at 2.1 bs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K,0)/1,000 sq. ft. 0.5# (K,0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9 Gandy 22 18 Lely 3 II 2 II Lesco (letter/numeric)E/15 D/14 PennMulch HVO G D ProScape SS G D	n off with water. nal use on turf, including golf con- train and the second s	be applied on dry or water dampened of ourses, sports fields, and landsd ate. Irming Jy after handling. his product. ng/eye protection/face protection. lical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. id water. doctor/physician if you feel unwell. ice/ attention.	concrete and should be removed from these are capes. Specific application rates and f Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local <b>Application Rates:</b> To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of proc SUGGESTED SPREADER S Spreader 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> 0.2 Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11 Gandy 22 Lely 3 II Lesco (letter/numeric)E/15 PennMulch HVO G ProScape SS G	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> l growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg duct per 100 m <sup>2</sup> . ETTINGS 24 kg K <sub>0</sub> /100 m <sup>2</sup> 1 <sup>1</sup> / <sub>2</sub> D 9 18 2 II D/14 D D
Application Rates:     To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K_0) or 100 sq. ft. apply this product at 4.2 lbs. per 1,000 sq. ft. apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K_0) per 1,000 sq. ft. apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre.     SUGGESTED SPREADER SETTINGS     Spreader   1½     Andressons AccuPro   F     Dearthway Rotary   11     Seandy   2     11   9     Gandy   22     11   9     Gandy   22     11   2     11   9     Gandy   22     11   2     11   2     11   2     11   2     11   2     11   2     11   2     11   2     11   2     11   11     11   2     11   1     11   2     11   1     11   2     11   1 <tr< td=""><td>n off with water. nal use on turf, including golf crosses and climate harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th Wear protective gloves/protective clothir If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and wast IF SWALLOWED: Call a POISON CENTER IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER or of If skin irritation occurs: Get medical advi If eye irritation persists: Get medical advi Store in a well-ventilated place.</td><td>be applied on dry or water dampened of ourses, sports fields, and landsd ate. Irming Jy after handling. his product. ng/eye protection/face protection. lical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. id water. doctor/physician if you feel unwell. ice/ attention.</td><td>concrete and should be removed from these are     capes. Specific application rates and f     Application Rates and Frequencies in Cana     professional landscape management plan     or area. A suggested application rate is 1 t     applied every 4 to 8 weeks during the local     Application Rates:     To feed at the rate of 0.24 kg Nitrogen (N)     Potash (K<sub>2</sub>O). Equivalent to 2.0 kg of proc     To feed at the rate of 0.12 kg Nitrogen (N)     Potash (K<sub>2</sub>O). Equivalent to 1.0 kg of proc     Spreader   0.48 kg K<sub>2</sub>O /100 m<sup>2</sup> 0.2     Lebanon Turf   2     Andersons AccuPro   F     Earthway Rotary   11     Gandy   22     Lely   3 II     Lesco (letter/numeric)E/15   PennMulch HVO     ProScape SS   G     Spyker   2</td><td>requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m<sup>2</sup> l growing season. or 0.48 kg duct per 100 m<sup>2</sup>. or 0.24 kg duct per 100 m<sup>2</sup>. ETTINGS 24 kg K<sub>2</sub>0 /100 m<sup>2</sup> 1<sup>1</sup>/<sub>2</sub> 9 18 2 II D/14 D D 1<sup>1</sup>/<sub>2</sub></td></tr<>	n off with water. nal use on turf, including golf crosses and climate harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th Wear protective gloves/protective clothir If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and wast IF SWALLOWED: Call a POISON CENTER IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER or of If skin irritation occurs: Get medical advi If eye irritation persists: Get medical advi Store in a well-ventilated place.	be applied on dry or water dampened of ourses, sports fields, and landsd ate. Irming Jy after handling. his product. ng/eye protection/face protection. lical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. id water. doctor/physician if you feel unwell. ice/ attention.	concrete and should be removed from these are     capes. Specific application rates and f     Application Rates and Frequencies in Cana     professional landscape management plan     or area. A suggested application rate is 1 t     applied every 4 to 8 weeks during the local     Application Rates:     To feed at the rate of 0.24 kg Nitrogen (N)     Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc     To feed at the rate of 0.12 kg Nitrogen (N)     Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of proc     Spreader   0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> 0.2     Lebanon Turf   2     Andersons AccuPro   F     Earthway Rotary   11     Gandy   22     Lely   3 II     Lesco (letter/numeric)E/15   PennMulch HVO     ProScape SS   G     Spyker   2	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> l growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg duct per 100 m <sup>2</sup> . ETTINGS 24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> 1 <sup>1</sup> / <sub>2</sub> 9 18 2 II D/14 D D 1 <sup>1</sup> / <sub>2</sub>
Application Rates:     To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) or 1,000 sq. ft. apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) or 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft. or 120 lbs. per acre.     SUGGESTED SPREADER SETTINGS     Spreader   1½     Andersons AccuPro   F     D   11   9     Gandy   22   18     Lely   3 II   2 II     Lely   3 II   2 II     PenoMulch HVO   G   D     ProScape SS   G   D     Spyker   2   1½	n off with water. nal use on turf, including golf con- total soils, regulations, and clima Hazard Statements Harmful if swallowed. Causes skin irritation. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th Wear protective gloves/protective clothir If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and wasf IF SWALLOWED: Call a POISON CENTEF IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER or of If skin irritation occurs: Get medical advi If eye irritation persists: Get medical advi Store in a well-ventilated place. Keep container tightly closed.	be applied on dry or water dampened of ourses, sports fields, and landso ate. Irning Jafter handling. his product. ng/eye protection/face protection. lical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. nd water. doctor/physician if you feel unwell. ice/ attention. rice/attention.	concrete and should be removed from these are capes. Specific application rates and f Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local <b>Application Rates:</b> To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of proc SUGGESTED SPREADER S Spreader 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> 0.2 Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11 Gandy 22 Lely 3 II Lesco (letter/numeric)E/15 PennMulch HVO G ProScape SS G Spyker 2 Vicon 17	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> I growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg duct per 100 m <sup>2</sup> . ETTINGS 24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> 1½ D 9 18 2 II D/14 D 1½ 14
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft. apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft. apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K <sub>2</sub> 0)/1,000 sq. ft. 0.5 # (K <sub>2</sub> 0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9 Gandy 22 18 Lely 3 II 2 II Lesco (letter/numeric)E/15 D/14 PennMulch HV0 G D ProScape SS G D Spyker 2 1½	n off with water. nal use on turf, including golf con- total soils, regulations, and climar Hazard Statements Harmful if swallowed. Causes skin irritation. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th Wear protective gloves/protective clothir If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and wasf IF SWALLOWED: Call a POISON CENTEF IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTEF or of Store in a well-ventilated place. Keep container tightly closed. Dispose of contents/container properly a	be applied on dry or water dampened of ourses, sports fields, and landso ate. Irning Jafter handling. his product. ng/eye protection/face protection. lical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. nd water. doctor/physician if you feel unwell. ice/ attention. rice/attention.	concrete and should be removed from these are capes. Specific application rates and f Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local <b>Application Rates:</b> To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of proc <b>SUGGESTED SPREADER S</b> Spreader 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> 0.2 Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11 Gandy 22 Lely 3 II Lesco (letter/numeric)E/15 PennMulch HVO G ProScape SS G Spyker 2 Vicon 17 These settings were calibrated and	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> I growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg duct per 100 m <sup>2</sup> . ETTINGS 24 kg K_0 /100 m <sup>2</sup> 1 $\frac{1}{2}$ D 9 18 2 II D/14 D D 1 $\frac{1}{2}$ 14 field tested.
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft. or 120 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft. apply this product at 2.1 lbs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K <sub>2</sub> 0)/1,000 sq. ft. 0.5 # (K <sub>2</sub> 0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9 Gandy 22 18 Lely 3 II 2 II Lesco (letter/numeric)E/15 D/14 PennMulch HVO G D ProScape SS G D Spyker 2 1½ These settings were calibrated and field tested.	n off with water. nal use on turf, including golf con- total soils, regulations, and clima Hazard Statements Harmful if swallowed. Causes skin Irritation. Causes eye irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th Wear protective gloves/protective clothir If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and wash IF SWALLOWED: Call a POISON CENTER IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER or of If skin irritation occurs: Get medical advi If eye irritation persists: Get medical advi Store locked up. Store in a well-ventilated place. Keep container tightly closed. Dispose of contents/container properly a Keep out of reach of children	be applied on dry or water dampened of ourses, sports fields, and landso ate. Irning Jafter handling. his product. ng/eye protection/face protection. lical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. nd water. doctor/physician if you feel unwell. ice/ attention. rice/attention.	concrete and should be removed from these are capes. Specific application rates and f Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local <b>Application Rates:</b> To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of proc SUGGESTED SPREADER S Spreader 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> 0.2 Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11 Gandy 22 Lely 3 II Lesco (letter/numeric)E/15 PennMulch HVO G ProScape SS G Spyker 2 Vicon 17	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> l growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg duct per 100 m <sup>2</sup> . ETTINGS 24 kg K <sub>0</sub> 0 /100 m <sup>2</sup> 1 $\frac{1}{2}$ D 9 18 2 II D/14 D 1 $\frac{1}{2}$ 14 field tested. ader, speed of
atomic intervent   atomic intervent     atomic intery intervent	n off with water. nal use on turf, including golf conserved as a solar of the second	be applied on dry or water dampened of ourses, sports fields, and landso ate. Urning Jy after handling. his product. ng/eye protection/face protection. lical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. ice/attention. ice/attention. ice/attention.	concrete and should be removed from these are capes. Specific application rates and f Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local <b>Application Rates:</b> To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of proc SUGGESTED SPREADER S Spreader 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> 0.2 Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11 Gandy 22 Lely 3 II Lesco (letter/numeric)E/15 PennMulch HVO G ProScape SS G Spyker 2 Vicon 17 These settings were calibrated and However, age and condition of sprea	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> l growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg duct per 100 m <sup>2</sup> . ETTINGS 24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> 1 $\frac{11}{2}$ D 18 2 II D/14 D D 1 $\frac{1}{2}$ 14 field tested. adder, speed of ay require slightly
ation of the set of th	n off with water. nal use on turf, including golf crosses and climate harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. Causes eye irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th Wear protective gloves/protective clothir If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and wast IF SWALLOWED: Call a POISON CENTER IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER or of If skin irritation occurs: Get medical advi If eye irritation persists: Get medical advi If eye irritation persists: Get medical advi Store in a well-ventilated place. Keep container tightly closed. Dispose of contents/container properly a Keep out of reach of children Precautionary Statements for disposal state and local regulations.	be applied on dry or water dampened of ourses, sports fields, and landso ate. Urning Jy after handling. his product. ng/eye protection/face protection. lical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. ice/attention. ice/attention. ice/attention.	concrete and should be removed from these are capes. Specific application rates and free professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local Application Rates: To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of proc SUGGESTED SPREADER S Spreader 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> 0.2 Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11 Gandy 22 Lely 3 II Lesco (letter/numeric)E/15 PennMulch HVO G ProScape SS G Spyker 2 Vicon 17 These settings were calibrated and However, age and condition of sprea	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> l growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg duct per 100 m <sup>2</sup> . ETTINGS 24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> 1 $\frac{11}{2}$ D 18 2 II D/14 D D 1 $\frac{1}{2}$ 14 field tested. adder, speed of ay require slightly
atom is product contains the micronutrient iron. Iron may stromptly after applications by sweeping or blowing. Do not wash ebanon brand fertilizers are formulated for profession etermined by the professional applicator based on lowing. The professional applicator based on lowing by the professional applicator based on lowing by the professional applicator based on lowing. The professional applicator based on lowing by the professional apply the profesis the professional apply the professic term a	n off with water. nal use on turf, including golf crosses and climate harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. Causes eye irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th Wear protective gloves/protective clothir If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and wast IF SWALLOWED: Call a POISON CENTER IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER or of If skin irritation occurs: Get medical advi If eye irritation persists: Get medical advi If eye irritation persists: Get medical advi Store in a well-ventilated place. Keep container tightly closed. Dispose of contents/container properly a Keep out of reach of children Precautionary Statements for disposal state and local regulations.	be applied on dry or water dampened of ourses, sports fields, and landso ate. Irning Iv after handling. his product. ng/eye protection/face protection. lical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. id water. doctor/physician if you feel unwell. ice/ attention. rice/attention. s required by law for waste pesticides. I - Dispose in accordance with all federal, s, nose, throat, and skin. In case of com	concrete and should be removed from these are capes. Specific application rates and free professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local Application Rates: To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of proc SUGGESTED SPREADER S Spreader 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> 0.2 Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11 Gandy 22 Lely 3 II Lesco (letter/numeric)E/15 PennMulch HVO G ProScape SS G Spyker 2 Vicon 17 These settings were calibrated and However, age and condition of spread operation and evenness of terrain m different settings for desired covera	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> l growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg duct per 100 m <sup>2</sup> . ETTINGS 24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> 1 $\frac{11}{2}$ D 18 2 II D/14 D D 1 $\frac{1}{2}$ 14 field tested. adder, speed of ay require slightly
Application Rates: For feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K_0) per 1,000 sq. ft. apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 1.5 lb. of Potash (K_0) per 1,000 sq. ft. apply this product at 2.1 bs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K_0)/1,000 sq. ft. 0.5 # (K_0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9 Gandy 22 18 Lely 3 II 2 II Lessco (letter/numeric)E/15 D/14 PronScape SS G D Spyker 2 1½ These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.	n off with water. nal use on turf, including golf crosses and climate harmful if swallowed. Causes skin irritation. Causes eye irritation. Causes eye irritation. Causes eye irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th Wear protective gloves/protective clothir If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and wash IF SWALLOWED: Call a POISON CENTER IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER or of If skin irritation occurs: Get medical advi If eye irritation persists: Get medical advi If eye irritation persists: Get medical advi Store in a well-ventilated place. Keep container tightly closed. Dispose of contents/container properly a Keep out of reach of children Precautionary Statements for disposal state and local regulations. Fertilizer may cause irritation of eyes	be applied on dry or water dampened of ourses, sports fields, and landso ate. Irning Iv after handling. his product. ng/eye protection/face protection. lical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. id water. doctor/physician if you feel unwell. ice/ attention. rice/attention. s required by law for waste pesticides. I - Dispose in accordance with all federal, s, nose, throat, and skin. In case of com	concrete and should be removed from these are capes. Specific application rates and free professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local Application Rates: To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of proc SUGGESTED SPREADER S Spreader 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> 0.2 Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11 Gandy 22 Lely 3 II Lesco (letter/numeric)E/15 PennMulch HVO G ProScape SS G Spyker 2 Vicon 17 These settings were calibrated and However, age and condition of spread operation and evenness of terrain m different settings for desired covera	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> l growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg duct per 100 m <sup>2</sup> . ETTINGS 24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> 1 $\frac{11}{2}$ D 18 2 II D/14 D D 1 $\frac{1}{2}$ 14 field tested. adder, speed of ay require slightly
otice: This product contains the micronutrient iron. Iron may st romptly after applications by sweeping or blowing. Do not wash ebanon brand fertilizers are formulated for profession etermined by the professional applicator based on lo Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 183 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 0.5 lb. of Potash (K <sub>2</sub> 0) per 1,000 sq. ft., apply this product at 2.1 bs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K <sub>2</sub> 0)/1,000 sq. ft. 0.5 # (K <sub>2</sub> 0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9 Gandy 22 18 Lely 3 II 2 II Lesco (letter/numeric)E/15 D/14 PennMulch HVO G D Syyker 2 1½ Vicon 17 14 These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.	n off with water. nal use on turf, including golf ca inclusional soils, regulations, and clima Harmful if swallowed. Causes skin irritation. Causes seye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th Wear protective gloves/protective clothin If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and wast IF SWALLOWED: Call a POISON CENTER IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER IF on SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER or of If skin irritation occurs: Get medical advi If eye irritation persists: Get medical advi Store in a well-ventilated place. Keep container tightly closed. Dispose of contents/container properly and Keep out of reach of children Precautionary Statements for disposal state and local regulations. Fertilizer may cause irritation of eyes skin or eyes, flush with plenty of wat	be applied on dry or water dampened of ourses, sports fields, and landso ate. Irning Iv after handling. his product. ng/eye protection/face protection. lical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. id water. doctor/physician if you feel unwell. ice/ attention. rice/attention. s required by law for waste pesticides. I - Dispose in accordance with all federal, s, nose, throat, and skin. In case of com	concrete and should be removed from these are capes. Specific application rates and f Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local <b>Application Rates:</b> To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of proc SUGGESTED SPREADER S Spreader 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> 0.2 Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11 Gandy 22 Lely 3 II Lessco (letter/numeric)E/15 PennMulch HVO G ProScape SS G Spyker 2 Vicon 17 These settings were calibrated and thowever, age and condition of spread operation and evenness of terrain m different settings for desired coverat tract with	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> l growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg duct per 100 m <sup>2</sup> . ETTINGS 24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> 1 $\frac{11}{2}$ D 1 $\frac{1}{2}$ D 1 $\frac{1}{2}$ 14 field tested. adder, speed of ay require slightly
ation of the set of th	n off with water. nal use on turf, including golf ca incal soils, regulations, and clima Hazard Statements Harmful if swallowed. Causes skin irritation. Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th Wear protective gloves/protective clothir If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and wash IF SWALLOWED: Call a POISON CENTER IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER or of if skin irritation occurs: Get medical advi If eye irritation persists: Get medical advi Store locked up. Store in a well-ventilated place. Keep container tightly closed. Dispose of contents/container properly and Keep out of reach of children Precautionary Statements for disposal state and local regulations. Fertilizer may cause irritation of eyes skin or eyes, flush with plenty of wath	be applied on dry or water dampened of ourses, sports fields, and landso ate. Irning Jafter handling. his product. ng/eye protection/face protection. lical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. ice/ attention. ice/ attention. ice/ attention. ice/ attention. s required by law for waste pesticides. I - Dispose in accordance with all federal, s, nose, throat, and skin. In case of com ter; for eyes get medical attention.	concrete and should be removed from these are capes. Specific application rates and f Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local Application Rates: To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of proc SUGGESTED SPREADER S Spreader 0.48 kg K <sub>2</sub> O/100 m <sup>2</sup> 0.2 Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11 Gandy 22 Lely 3 II Lesco (letter/numeric)E/15 PennMulch HVO G ProScape SS G Spyker 2 Vicon 17 These settings were calibrated and However, age and condition of spread operation and evenness of terrain m different settings for desired coveration that with	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> l growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg duct per 100 m <sup>2</sup> . ETTINGS 24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> 1 $\frac{11}{2}$ D 18 2 II D/14 D D 1 $\frac{1}{2}$ 14 field tested. adder, speed of ay require slightly
Application Rates: To feed at the rate of 0.5 lb. Nitrogen (N) or 1 lb. of Potash (K_0) ter 1,000 sq. ft., apply this product at 4.2 lbs. per 1,000 sq. ft. or 83 lbs. per acre. To feed at the rate of 0.25 lb. Nitrogen (N) or 1.5 lb. of Potash (K_0) per 1,000 sq. ft., apply this product at 2.1 bs. per 1,000 sq. ft. or 120 lbs. per acre. SUGGESTED SPREADER SETTINGS Spreader 1# (K_0)/1,000 sq. ft. 0.5 # (K_0)/1,000 sq. ft. Lebanon Turf 2 1½ Andersons AccuPro F D Earthway Rotary 11 9 Gandy 22 18 Lely 3 II 2 II Lesco (letter/numeric)E/15 D/14 PennMulch HVO G D ProScape SS G D Spyker 2 1½ These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.	n off with water. nal use on turf, including golf con- tract soils, regulations, and clima Hazard Statements Harmful if swallowed. Causes skin irritation. Causes sein irritation. Causes eye irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation. Precautionary Statements for handling Wash hands and exposed skin thorough Do not eat, drink or smoke when using th Wear protective gloves/protective clothir If in eyes, rinse with water and seek med Rinse mouth. Take off contaminated clothing and wash IF SWALLOWED: Call a POISON CENTER IF ON SKIN: Wash with plenty of soap an IF INHALED: Call » POISON CENTER or of If skin irritation occurs: Get medical advi If eye irritation persists: Get medical advi Store locked up. Store in a well-ventilated place. Keep container tightly closed. Dispose of contents/container properly at Keep out of reach of children Precautionary Statements for disposal state and local regulations. Fertilizer may cause irritation of eyes skin or eyes, flush with plenty of wat Use in accordance with recommenda such as, but not limited to a certified	be applied on dry or water dampened of ourses, sports fields, and landso ate. Irning Jaly after handling. his product. ng/eye protection/face protection. lical attention if symptoms persist. h before reuse. R or doctor/physician if you feel unwell. ice/ attention. ice/ attention.	concrete and should be removed from these are capes. Specific application rates and f Application Rates and Frequencies in Cana professional landscape management plan or area. A suggested application rate is 1 t applied every 4 to 8 weeks during the local Application Rates: To feed at the rate of 0.24 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 2.0 kg of proc To feed at the rate of 0.12 kg Nitrogen (N) Potash (K <sub>2</sub> O). Equivalent to 1.0 kg of proc SUGGESTED SPREADER S Spreader 0.48 kg K <sub>2</sub> O /100 m <sup>2</sup> 0.2 Lebanon Turf 2 Andersons AccuPro F Earthway Rotary 11 Gandy 22 Lely 3 II Lesco (letter/numeric)E/15 PennMulch HVO G ProScape SS G Spyker 2 Vicon 17 These settings were calibrated and t However, age and condition of spreat operation and evenness of terrain m different settings for desired coverat that with	requencies are ada: Follow your for your local cours to 2.8 Kg for 100 m <sup>2</sup> l growing season. or 0.48 kg duct per 100 m <sup>2</sup> . or 0.24 kg duct per 100 m <sup>2</sup> . ETTINGS 24 kg K <sub>2</sub> 0 /100 m <sup>2</sup> 1 $\frac{11}{2}$ D 18 2 II D/14 D D 1 $\frac{1}{2}$ 14 field tested. adder, speed of ay require slightly

Do not apply more than 3.2 lbs. total nitrogen (N) per year.

Do not apply to frozen turf.

on impervious surfaces back into your turf.

nutrient management plan.

WARRANTY NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your turf, and sweep any product that lands

For technical assistance or more information about our products visit www.LebanonTurf.com

### Lebanon Seaboard Corporation

1600 E. Cumberland St • Lebanon, PA 17042

800-233-0628 • 717-273-1685

### 40 Lbs. (18.14 kg) Net Weight