

# 19-0-5

## .37% Prodiamine

### GUARANTEED ANALYSIS

Total Nitrogen (N) .....	19.0%
19.0% Urea nitrogen*	
Soluble Potash (K <sub>2</sub> O) .....	5.0%

Plant foods derived from: Urea and Muriate of Potash.

495-5100

polypocketsm



## PROFESSIONAL TURF FERTILIZER

### PRODIAMINE 0.37 PLUS

FOR PRE-EMERGENCE CONTROL OF GRASS AND BROADLEAF WEEDS IN LAWNS AND ORNAMENTAL TURF

(excluding golf course putting greens)

LANDSCAPE ORNAMENTALS, ESTABLISHED PERENNIALS AND WILDFLOWER PLANTINGS

STOPS MANY WEED GRASSES BEFORE THEY GROW

PREVENTS CRABGRASS, ANNUAL BLUEGRASS, FOXTAIL

#### ACTIVE INGREDIENT:

Prodiamine (CAS # 29091-21-2) ..... 0.37%

INERT INGREDIENTS: ..... 99.63%

TOTAL: ..... 100.00%

EPA Reg. No.53883-172-57131

EPA Est. No. 57131-WI-1

## Net Wt. 50 lbs.

### KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

#### FIRST AID

<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>Call a poison control center or physician immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by the poison control center or physician.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.	

#### ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is a pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens), landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds, germination and root development. When used according to directions, it will control; barnyardgrass; bluegrass; annual (Poa annua); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass<sup>1</sup>; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass<sup>3</sup>, shepherd's purse<sup>2</sup>, signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).


<sup>1</sup>In many areas a single application of 4 to 9.3 pounds/1,000 square feet (175 to 405 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 4 to 6.2 pounds/1,000 square feet (175 to 270 pounds/acre), followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).

<sup>2</sup>For this weed, apply in late summer, fall or winter prior to germination.

<sup>3</sup>Suppression only.

**WHEN TO APPLY TO ESTABLISHED TURF:** This product must be applied prior to germination of weed seeds, as it will not control established weeds.

#### THIS MAP SHOWS APPROXIMATE CRABGRASS GERMINATION DATES



**APPLICATION DIRECTIONS:** Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days application.

**USE PRECAUTIONS:** The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil). Do not apply to newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control. Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (Poa annua) are desirable species. Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.

**RATES OF APPLICATION:** This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

MINIMUM AND MAXIMUM APPLICATION RATES (by Turf Grass Species)		
Turf Species	Minimum Rate	Maximum Annual Rate <sup>1</sup>
Bahiagrass, Bermudagrass <sup>2</sup> Centipedegrass, Zoysiagrass, Seashore Paspalum, St. Augustinegrass Tall Fescue (including turf-type)	4 lbs/1,000 sq. ft. (175 lbs/acre)	9.3 lbs/1,000 sq. ft. per year (405 lbs/acre per year)
Buffalograss Kentucky Bluegrass Perennial Ryegrass	4 lbs/1,000 sq. ft. (175 lbs/acre)	6.2 lbs/1,000 sq. ft. per year (270 lbs/acre per year)
Creeping Red Fescue	4 lbs/1,000 sq. ft. (175 lbs/acre)	4.6 lbs/1,000 sq. ft. per year (202 lbs/acre per year)
Creeping Bentgrass	4 lbs/1,000 sq. ft. (175 lbs/acre)	4.0 lbs/1,000 sq. ft. per year (175 lbs/acre per year)

<sup>1</sup>These are the maximum rates per calendar year.

<sup>2</sup>May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 3.3 pounds/1,000 square feet (145 pounds/acre). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

**WHEN TO OVERSEED AFTER APPLICATION:** This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/reseeding results (see map above).

USE RATE	MONTHS BEFORE OVERSEEDING		
	North	Transition	South
3 pounds/1,000 sq. ft. (135 pounds/acre)	4	4	4
4 pounds/1,000 sq. ft. (175 pounds/acre)	5	4	4
4.6 pounds/1,000 sq. ft. (202 pounds/acre)	6	5	5
4.9 pounds/1,000 sq. ft. (215 pounds/acre)	-	6	6
6.2 pounds/1,000 sq. ft. (270 pounds/acre)	-	7	7
7 pounds/1,000 sq. ft. (308 pounds/acre)	-	-	9
8 pounds/1,000 sq. ft. (350 pounds/acre)	-	-	10
9.3 pounds/1,000 sq. ft. (405 pounds/acre)	-	-	12

**LANDSCAPE ORNAMENTALS:** (including established perennials and wildflower plantings): This product may be applied for residual pre-emergence weed control in ornamentals. Apply at the rate of 4 to 9.3 pounds/1,000 square feet (175 to 405 pounds/acre) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 9.3 pounds/1,000 square feet (405 pounds/acre) per year.

**APPLICATION TIMING AND INFORMATION:** This product may be applied to newly transplanted and established ornamentals a broadcast, over the top or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grfts have taken to avoid any inhibition of the tissue union. This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate post-emergence herbicide.

**TOLERANT ORNAMENTAL SPECIES:** This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

#### SCIENTIFIC NAME

Abies spp.  
Abelia grandiflora  
Acer palmatum  
A. Platanoides  
Achillea spp.  
Achillea chinensis  
Agapanthus africanus  
Agapanthus orientalis  
Akaihaq umbata  
Allium cernuum  
Anemone hybrida  
Aquilegia spp.  
Arocladophytos densiflora  
Ariocnece calendula  
Ariemesia spp.  
Aster spp.  
Aster X Frikarti  
Athinum Fliz-Jemina  
Aucuba japonica  
Begonia spp.  
Begonia glabrymensis  
B. Julianae  
B. membranis  
B. Thunbergii  
B. Veruculosa  
Bergenia cordifolia  
Bolonia asteroides  
Buddleia davidii  
Buxus microphylla  
Callistemon viminalis  
Calluna vulgaris  
Campanula carpatica  
Campis X Tagliacara  
Carpobrotus seduis  
Cassia artemisoides  
Ceanothus rigidus  
Ceratostigma plumbagionoides  
Chamaecyparis pisifera  
Cleyera japonica  
Citrus spp.  
Conopsis spp.  
Cornus florida  
C. Stolonifera  
Cortaderia seloana  
Cotoneaster apiculatus  
C. buxifolius  
C. dammeri  
C. Microphyllus  
Cretaceous spp.  
Crocosmia spp.  
Cupressus sempervirens  
Dactyloctenium aegyptium  
Dactylis glomerata  
Daphnium spp.  
Dianthus deltoides  
D. griffithii  
Dodonaea viscosa  
Echinacea purpurea  
Eleagnus pungens  
Eunymus fortunei  
E. japonica  
E. kautschovicia  
Falsa japonica  
Forsythia intermedia  
Forsythia suspensa  
F. viridissima  
Gallardia spp.  
Gardenia jasminoides  
Gaura spp.  
Gentiana dahurica  
Geranium cinereum  
Glaucolobus spp.  
Gypsophila repens  
Hedera helix  
Helleborus spp.  
Hemerocallis spp.  
Hibiscus  
Hibiscus spp.  
Hibiscus rosa-sinensis  
Hosta plantaginea  
H. Strobiliana  
Houttuynia cordata var. variegata  
Hydrangea macrophylla  
Ilex Berni  
Ilex cornuta  
I. crenata  
I. opaca  
I. vomitoria  
Iris ensata  
Iris sibirica  
Iris spp.  
Jasminum nudiflorum  
Juniperus chinensis  
J. conferta  
J. davurica  
J. horizontalis  
Juglans spp.  
Justicia brandegeana  
Lagerstromia indica  
Lagestromia indica X Faulrei  
Lantana montevidensis  
Lavender spp.  
Leonopodium alpinum  
Ligustrum amurense  
Ligustrum sinense  
L. japonicum  
L. lucidum  
Lilium spp.  
Liriope muscari  
Liriope muscari var. variegata  
Liriope spicata  
Lobelia Cardinalis  
Lonicera japonica  
L. latianca  
Lorapetalum chinense  
Lythrum spp.  
Magnolia spp.  
Malaeophora tubola  
Malus spp.  
Miscanthus sinensis  
Nandina domestica  
Narcissus spp.  
Nerium oleander  
Olea europaea  
Ophiopogon japonicus  
Omanthus heterophyllus  
Ostrya serratum  
Ostrya serratum  
Oxydendrum abotocum  
Paeonia suffruticosa  
Persea americana  
Perovskia atriplicifolia  
Persea americana  
Phloxia Fraseri  
Physlogia Virginiana  
Pilea spp.  
Pieris japonica  
Pinus brutia  
P. cananensis  
P. Elliptica  
P. halepensis  
P. nigra  
P. palustris

COMMON NAME  
Fl. species\*\* (Balsam, Frasier, Noble, etc.)  
Abelia: Sherwood  
Japanese Maple  
Norway Maple  
Yarrow: King Edward  
Kiwi  
Lily-of-the-Nile (African Lily)

Five-leaf or Chocolate Vine  
Lady's, Leak, Nodding Onion  
Japanese Anemone  
Aquilegia: Red and Gold  
Vine Hill Marzanita  
Cape Weed  
Wormwood: Silver Mound, Castle  
Aster: Bonnie Blue, Purple Dome

Lady Fern; Fern Lady  
Japanese Aucuba  
Ebrous Begonia; Hardy Grandis  
Barbery  
Wintergreen Barbery  
Mentor Barbery  
Japanese Barbery  
Warty Barbery

Shovbank  
Butterfly-Bush (Dwarf Blue); Royal Red  
Japanese Boxwood  
Weeping Bottlebrush  
Scottish Heather  
Tussock Beilflower (White Clips)  
Trumpet Flower; Madame Galen  
Fotemort Fig (Ice Plant)  
Feathery Cassia  
Wild Lic

False Cypress  
Cleyera  
Citrus species\*  
Conopsis (Calliopsis) Early Sunrise, Moonbeam  
Flowering Dogwood  
American Dogwood  
Pampas Grass  
Cranberry Cotoneaster  
Cotoneaster  
Bearberry Cotoneaster  
Rockspray Cotoneaster  
Hawthorne  
Lucifer  
Italian Cypress  
White Trailing Ice Plant  
Cooper; Pink  
Larkspur; Blue Elf  
Dianthus; Maiden Pinks "Zing"  
Cherry Pink  
Hop Bitch  
Coneflower; Purple; Magnus  
Silverberry  
Wintercreeper  
Evergreen Euonymus  
Spreading Euonymus  
Japanese Aralia  
Border Forsythia  
Weeping Forsythia  
Greenstein; Forsythia  
Gallardia, Blanket Flower; "Goblin"  
Gardenia, Cape-Jasmine

Gentian  
Cranesbill  
Glaucolobus species\*\*  
Baby's Breathe  
English Ivy  
Succesa  
Daylily; Aztec Gold; Stella de Oro, Tender Love  
Rose of Sharon\*\*  
Mallow; Disco Belle White  
Chinese Hibiscus  
Hosta  
Hosta, Seersucker\*

Boleif Hydrangea  
Holy  
Chinese Holly  
Japanese Holly  
American Holly  
Raupon Holly; Schilling's  
Sword-Leaved Iris; Jodlesong  
Siberian Iris; Cabernet  
Iris species\*\*  
Winter Jasmine  
Chinese Juniper; Nick's Compact; Parsomni  
Shore Juniper; Blue Pacific  
Parsoni  
Creeping Juniper  
Wainut\*

Stump Plant  
Cape Myrtle  
Crape Myrtle; Tuscarora  
Weeping Lantana  
Lavender; Munstead  
Edelweiss  
Anur Privet  
Chinese Privet; Variegata  
Japanese Privet  
Glossy Privet (Waxleaf)

Lily-Jazz  
Big Blue Lily; Silver Mound, Evergreen Giant  
Liriope, Variegated  
Liriope, Creeping  
Cardinal Flower, Indian Pink  
Japanese Honeysuckle  
Tatarian Honeysuckle  
Burgundy  
Loosestrife; Modern Pink  
Magnolia species\*\*  
Ice Plant\*  
Crabapple  
Yaku Jima; Silberfeder  
Heavenly Bamboo  
Narcissus species\*  
Oleander  
Olive\*  
Mondo Grass  
Osmanthus (False Holly); Gulf Tite  
Trailing African Daisy  
Sourwood  
Tree Peony  
Fountain Grass (Dwarf)

Avocado\*  
Fraser's Phloxia (Redlip)  
Dragonhead, False Vivid  
Spruce species\*\* (Colorado Blue, Norway, etc.)  
Lily-of-the-Valley Bush  
Calabrian Pine  
Canary Island Pine  
Slash Pine  
Aleppo Pine  
Austrian Pine  
Longleaf Pine

P. radiata  
P. strobus  
P. sylvestris  
P. taeda  
P. Thunbergiana  
P. Virginiana  
Pistachio spp.  
Pittosporum rhombifolium  
P. tobira  
Podocarpus macrophyllus  
P. umus spp.  
P. taeda  
Pseudotsuga menziesii  
Pyracantha coccinea  
P. fortuneana  
P. koidzumii  
Pyrus spp.  
Quercus shumardii  
Quercus spp.  
Raphiolepis indica  
Raphiolepis umbellata  
Rhododendron  
(including Azalea)

Rosa banksiae  
Rosmarinus officinalis  
Rudbeckia spp.  
Santolina virens  
Saxifraga spp.  
Scaevola spp.  
Sedum album  
Sedum cauticola  
Sedum dasycyllum  
Sedum spuriun  
Spiraea bumalda  
Syringum paniculatum  
T. media  
T. nigra  
Teucrium spp.  
Thalictrum dipercarpum  
Thuja occidentalis  
Trachelospermum asiaticum  
Tsuga canadensis  
Tulipa spp.  
Veronica spp.  
Viburnum japonicum  
V. odoratissimum  
V. plicatum  
V. rigratum  
V. suspensum  
V. tinus  
V. trilobum  
V. wrightii  
Vinca major  
Vinca minor  
Vitis spp.  
Weigela florida  
Yucca aloifolia  
Y. filifemosa  
\*\*Not for use on container grown plants.

Monterey Pine  
Eastern White Pine  
Scotch Pine  
Loblolly Pine  
Japanese Black Pine  
Virginia Pine  
Pistachio  
Queensland Pittosporum  
Japanese Pittosporum  
Japanese Yew  
English Laurel  
Almond, Apricot, Nectarine, Peach, Plum and Plum\*  
Douglas Fir\*\*  
Fretform (Scotiae)  
Fretform  
Bradford Pear spp.  
Oak, Shumard's Red  
Oak species  
Indian Hawthorne  
Yeda Hawthorne  
Coral Balls; \*Famosa; \*\*Hiro-crimson; \*PJM  
Delaware Valley White; \*Roseum Elegans;  
Flame Creeper; \*Grand Crimson; \*Wakelabisu  
White Gumbo; \*George L. Tabor  
Lady Banks's Rose  
Rosenary

Black-Eyed Susan; Goldstium  
Santolina  
Saxifrage; Purple Dome  
Pin cushion Flower  
Stonecrop  
Stonecrop; Lidakense  
Stonecrop  
Spiraea; Dragon's Blood  
Spiraea; Anthony Waterer  
Austrian Blushcherry  
Japanese Yew  
Yew  
Germander  
Meadow Rue  
American Arbutovae  
Star Jasmine  
Canada Hemlock  
Tulip species  
Veronica; Speedwell; Sunny Border  
Japanese Viburnum  
Sweet Viburnum  
Japanese Snowball  
Canary Island Viburnum  
Arrowood Viburnum  
Laurusinus  
Cranberry Bush  
Leatherleaf Viburnum  
Greater Penwinkle; Vinca  
Common Penwinkle; Vinca  
Grape\*  
Old Fashioned Weigela  
Spanish Bayonet; Yucca  
Adam's Needle; Yucca

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.  
**Pesticide Storage:** Store in original container in a dry place. Store in locked area out of reach of children and domestic animals.  
**Pesticide Disposal:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinseate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

**Container Disposal:** Non-refillable container. Do not reuse or refill this container. Empty container through use according to label instructions. Do not rinse unless required for recycling. Offer for recycling, if available. If recycling is not available, then dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

#### CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of The Manufacturer or the Seller. All such risks shall be assumed by the Buyer.

The Manufacturer warrants that this product conforms to the chemical description on the label, and is reasonably fit for the purposes set forth in the DIRECTIONS FOR USE, when it is used in accordance with such directions, subject to the inherent risks mentioned above.

THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THIS WARRANTY EXTENDS TO AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS; WARNING, OR CAUTIONS. BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED AT THE MANUFACTURER'S OPTION TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. THE MANUFACTURER and the seller offer this product, and the Buyer and User accept it, subject to the foregoing CONDITIONS OF SALE AND WARRANTY.

# EC GROW

Distributed By Eau Claire Co-op Oil Co., P.O. Box 837, Eau Claire, WI 54702