

Edgeless® Liquid

Turf Growth Regulator

SPECIMEN



For edging and banding applications to control growth and reduce trimming requirements of perennial turfgrass species.

Active Ingredients

flurprimidol: α -(1-methylethyl)- α -[4-(trifluoromethoxy)phenyl]-5-pyrimidinemethanol 13.26%
trinexapac-ethyl: 4-(cyclopropyl-alpha-hydroxymethylene)-3,5-dioxo-
cyclohexanecarboxylic acid ethyl ester..... 5.00%

Other Ingredients 81.74%

TOTAL 100.00%

Contains 1.10 pounds of flurprimidol per gallon of product.
Contains 0.41 pound of trinexapac-ethyl per gallon of product.

Keep Out of Reach of Children

WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Refer to the inside of label booklet for additional precautionary information and Directions for Use including First Aid and Storage and Disposal.

NOTICE: Read the entire label before using. Use only according to label directions. Before buying or using this product, read *Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies* inside label booklet.

EPA Reg. No. 67690-46
FPL20110926

Manufactured for: SePRO Corporation 11550 North Meridian Street, Suite 600
Carmel, IN 46032, U.S.A.

SPECIALTY CHEMICAL: Do not ship or store with food, feeds, drugs or clothing.

Refer to inside of label booklet for additional precautionary information and directions for use, including first aid and storage and disposal.

Notice: Read the entire label before using. Use only according to label directions. Before buying or using this product, read *Terms and Conditions of Use, Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies* at the end of the label booklet. If terms are unacceptable, return at once unopened.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

WARNING / AVISO

Causes substantial but temporary eye injury. Do not get in eyes or on skin or clothing. Harmful if swallowed or inhaled. Avoid breathing spray mist. Wear protective eyewear and chemical-resistant gloves.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants;
- Shoes plus socks;
- Chemical-resistant gloves (such as nitrile, butyl rubber, neoprene, or barrier laminate); and
- Protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency endangering health or the environment involving this product, call **INFOTRAC at 1-800-535-5053**.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Ground Water Advisory

This pesticide has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product is classified as having potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs, will reduce the potential loading of flurprimidol from runoff and sediment.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read all directions for use carefully before applying. Use only according to label directions.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170).

The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated area without footwear until sprays have dried.

Avoiding Injurious Spray Drift

Applications should be made only when there is little or no hazard for spray drift. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants. Do not spray when wind is blowing toward desirable susceptible crops or ornamental plants near enough to be injured. Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions.

Product Information for Growth Regulation of Perennial Turfgrasses

Edgeless integrates patented[†] synergy of turf growth regulator technology for use on both warm and cool-season perennial turfgrasses on golf courses. Edgeless reduces stem elongation and leaf blade length in perennial turfgrasses resulting in a more compact and dense growth form. Growth regulation from Edgeless results from suppression of the plant hormone, gibberellic acid (GA), responsible for cell elongation in most plants.

[†]The synergy derived from the combination of Type II Class A and Type II Class B PGRs is protected by United States Patent No. 7,135,435. Additional patent rights pending.

Broadcast treatments should be made on medium to high quality turfgrass areas. An appropriate fertility program for the desired turf species should be followed in conjunction with Edgeless applications to provide the best turfgrass enhancement and reduce potential for discoloration.

NOTICE TO USER: Turfgrass responses to Edgeless may vary within turfgrass species due to the large number of cultivars and varieties available. Neither the manufacturer nor seller has determined if Edgeless can be used safely or effectively on turfgrass species not mentioned on this label. For turfgrass species not listed on this label the user should apply Edgeless to a small test area to determine growth response and desired level of growth regulation prior to large scale applications.

Use Precautions for Perennial Turfgrasses

- **Edgeless is not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production or for research purposes.**
- Do not apply to shrubs, bedding plants, and/or food plants.
- Not for use on turfgrasses under stress due to temperature and moisture extremes and disease, and/or insect pressures.
- Applications of Edgeless to newly seeded turfgrasses should be delayed until turf is well established and actively growing.
- Edgeless should not be applied until 6 to 8 weeks after turfgrass sprigging or laying sod. Turfgrass should be well established and actively growing prior to application.
- Do not apply to saturated soils or when a significant moisture event is anticipated.
- Do not apply to turf used for livestock production.
- Multiple applications of Edgeless can be made yearly, but do not apply more than 350 fl oz/A of Edgeless per acre per year.
- **Chemigation:** Do not apply Edgeless through any type of irrigation system.

Application Timing

Applications of Edgeless should be made to turfgrass actively growing. Spring applications should be made after resumption of active seasonal growth of turfgrass. The final application of the season should be timed a minimum of 4 weeks before the onset of inactive grass growth or winter dormancy. Applications to overseeded turfgrasses in dormant bermudagrass stands should be made 4 weeks prior to expected bermudagrass green-up.

Irrigation

Rainfall or irrigation should be delayed at least 2 hours after application or until product has dried on the leaf surface but should occur within 24 hours after application. Avoid mowing turfgrass areas treated with Edgeless until after rainfall or irrigation occurs.

Turf Color and Post Treatment Turf Management

Turfgrass treated with Edgeless may appear darker green in color. This color change, which appears 1 to 2 weeks after treatment, may persist an additional 3 to 6 weeks. Treated areas should be managed to encourage the growth of a healthy vigorous turf.

Application Directions

Mixing Directions

Add Edgeless to a spray tank half filled with clean water while agitating. Allow sufficient mixing time to ensure consistent mixing of Edgeless. Finish filling the spray tank. Continue agitation throughout the spraying

operation to ensure uniform application. The use of a coloring agent or foam to mark areas already sprayed is suggested for uniform application without skips and overlaps.

EDGING AND BANDING APPLICATIONS FOR GROWTH REGULATION OF PERENNIAL TURFGRASS SPECIES

Edgeless can be applied to turfgrass in edging and banding applications along the perimeter of lawns, landscape beds, sidewalks, curbs, parking lots, driveways, posts, mailboxes, building structures, gravestones, and fences. Edgeless must be applied in a 6 inch band width with a single nozzle sprayer. Repeat at 8 to 12 week intervals or as needed for turf growth regulation.

TABLE 1	
Edgeless Edging/Banding Rates for Growth Regulation of Perennial Turfgrass	
Turfgrass Species	Fl. Oz. Edgeless/A
Cool Season Turfgrasses	
Bentgrass	30 - 60
Kentucky Bluegrass	40 - 80
Perennial Ryegrass	40 - 80
Tall Fescue	40 - 80
Warm Season Turfgrasses	
328 Hybrid Bermudagrass	20 - 40
419 Hybrid Bermudagrass	30 - 60
Centipedegrass	20 - 40
Common Bermudagrass	40 - 80
Seashore Paspalum	30 - 60
St. Augustinegrass	30 - 60
Zoysiagrass	30 - 60

In order to deliver the correct rate of Edgeless to the desired turf species, users should properly calibrate their sprayer to determine spray volume in gallons per acre. Following proper calibration, use Table 2 to determine fluid ounces of Edgeless per 1 gallon of water required to apply the targeted rate found in Table 1 at various spray volumes.

Table 2							
Fluid Ounces of Edgeless Per Gallon of Water Required to Apply Targeted Rates of Edgeless at Various Spray Volumes.							
Target Rate of Edgeless (Fl oz./A)	Gallon Per Acre (GPA) Spray Volume						
	20 GPA	30 GPA	40 GPA	50 GPA	60 GPA	70 GPA	80 GPA
20	1.0	0.7	0.5	0.4	0.3	0.3	0.3
30	1.5	1.0	0.8	0.6	0.5	0.4	0.4
40	2.0	1.3	1.0	0.8	0.7	0.6	0.5
50	2.5	1.7	1.3	1.0	0.8	0.7	0.6
60	3.0	2.0	1.5	1.2	1.0	0.9	0.8
70	3.5	2.3	1.8	1.4	1.2	1.0	0.9
80	4.0	2.7	2.0	1.6	1.3	1.1	1.0

EQUIPMENT CALIBRATION FOR EDGING AND BANDING APPLICATIONS

Proper application rate, volume and placement are important to ensure efficacy with Edgeless. SePRO recommends specific application equipment and spray techniques to maximize efficacy of Edgeless. **All spray equipment must be properly calibrated before applying Edgeless.** For optimum application, and to provide the best results, follow the 3 steps below:

1. **Determine Desired Rate (Fl. Oz./A) Based upon Targeted Turfgrass Species.** Use Table 1 to identify the proper use rate (use higher rates when environmental conditions favor vigorous growth of turfgrass species and when longer regulation is desired).
 - a. For example, apply 40 to 80 ounces per acre to effectively regulate Kentucky Bluegrass.
2. **Properly Calibrate Sprayer in Gallons Per Acre (GPA).** Follow instructions below for specific calibration recommendations to determine GPA for single nozzle sprayers.

- a. **Nozzle** – An even distribution nozzle is important for uniform coverage and resulting growth regulation. **SePRO recommends a TeeJet® 8002E nozzle for applications of Edgeless.**
- b. **Pressure (PSI) at Nozzle** – Maintaining a consistent pressure at the spray nozzle is difficult with a single nozzle sprayer when using a backpack or other hand-held compression sprayers. In order to maintain a consistent pressure at the nozzle, SePRO recommends using a pressure regulating device to target approximately 20 PSI when using the 8002E nozzle. **To target approximately 20 PSI, SePRO recommends the Chapin CFValve™ (Model CFV1X-R3-8BSP) designed to maintain 21 PSI.**
- **Sprayer/Nozzle Construction:** Remove the nozzle cap and nozzle and replace with the 8002E nozzle. Next, thread the Chapin CFValve between the nozzle cap and the end of spray wand. The CFValve is designed to fit a 3/8" thread. Larger adapters (11/16") can be ordered from Chapin (www.chapinmfg.com) if needed.
- c. **Height of Spray Tip** – Edgeless must be applied at a 6 inch wide band with a single nozzle sprayer. **SePRO recommends the 8002E nozzle which will apply Edgeless in a 6 inch band when held at 3.5 inches above the canopy.** Some users may design a guide that extends 3.5 inches below the spray tip to better target the proper height.
- d. **Walking Speed** – Edgeless is to be applied while walking at a consistent speed and holding the boom steady over the turf surface. It must NOT be applied by moving the spray wand back and forth over an area – this will result in non-uniform regulation. **Edgeless should be applied at 2, 2.5 or 3.0 mph.**
- To calibrate your walking speed, determine **how long in seconds it takes to walk 100 feet or 33 yards.** Use Table 3 to determine your walking speed and resulting Gallons Per Acre (GPA). Additionally, Table 3 shows the approximate linear feet that could be treated per gallon of spray solution (assumes 6-inch wide spray band).

Walking Speed (mph)	Time (seconds) to walk 100 ft	Gallons Per Acre*	Treatable Linear Feet Per Gallon Spray Solution ¹
2.0	34	77	1,131 feet
2.5	27	62	1,405 feet
3.0	23	51	1,708 feet

¹ Assumes a 6-inch wide spray band and utilizing the TeeJet 8002E nozzle applied at approximately 20 PSI

* For single nozzle sprayers calibrated at different GPA than shown below, refer to Table 2 for more dilution specifications

3. Dilute your Spray Solution Based upon Walking Speed and Desired Use Rate. Use Table 4 (dilution chart) to determine fluid ounces of Edgeless to mix per gallon of water based upon walking speed and targeted spray rate. Use Table 1 (Edgeless Edging/Banding Rates for Growth Regulation of Perennial Turfgrass) to determine spray rate for a specific turf species.

Walking Speed (mph)	GPA ²	Edgeless Rate (fl oz/A) (see <i>Edgeless Rates in Table 1</i>)						
		20	30	40	50	60	70	80
2	77	0.26	0.39	0.52	0.65	0.78	0.91	1.04
2.5	62	0.32	0.48	0.65	0.81	0.97	1.13	1.29
3	51	0.39	0.59	0.78	0.98	1.18	1.37	1.57

¹ Assumes using the TeeJet 8002E Nozzle applied at approximately 20 PSI and a 6-inch wide spray band for all calculations.

² For single nozzle sprayers calibrated at different GPA than shown below, refer to Table 2 for more dilution specifications.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

Pesticide Storage

Store in original container only. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

Container Handling

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Triple rinse containers small enough to shake (capacity ≤ 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Triple rinse containers too large to shake (capacity > 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable Container. Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

Triple rinse as follows: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

When this container is empty, replace the cap and seal all openings that have been opened during use; return the container to the point of purchase or to a designated location. This container must only be refilled with a pesticide product. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transport. **DO NOT** transport if this container is damaged or leaking. If the container is damaged, or leaking, or obsolete and not returned to the point of purchase or to a designated location, triple rinse emptied container and offer for recycling, if available, or dispose of container in compliance with state and local regulations.

TERMS AND CONDITIONS OF USE

If terms of the following *Warranty Disclaimer*, *Inherent Risks of Use* and *Limitation of Remedies* are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, to the extent consistent with applicable law, use by the buyer or any other user constitutes acceptance of the terms under *Warranty Disclaimer*, *Inherent Risks of Use*, and *Limitation of Remedies*.

WARRANTY DISCLAIMER

SePRO Corporation warrants that the product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SEPRO CORPORATION MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of SePRO Corporation or the seller. To the extent consistent with applicable law, all such risks shall be assumed by buyer.

LIMITATION OF REMEDIES

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories) shall be limited to, at SePRO Corporation's election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

To the extent consistent with applicable law, SePRO Corporation shall not be liable for losses or damages resulting from handling or use of this product unless SePRO Corporation is promptly notified of such losses or damages in writing. In no case shall SePRO Corporation be liable for consequential or incidental damages or losses.

The terms of the *Warranty Disclaimer*, *Inherent Risks of Use* and this *Limitation of Remedies* cannot be varied by any written or verbal statements or agreements. No employee or sales agent of SePRO Corporation or the seller is authorized to vary or exceed the terms of the *Warranty Disclaimer* or this *Limitation of Remedies* in any manner.

© Copyright 2013 SePRO Corporation

* Trademark of SePRO Corporation

TeeJet is a registered trademark of Spraying Systems Co.

CFValve™ is a trademark of GATE LLC.



SePRO Corporation
11550 North Meridian Street, Suite 600
Carmel, IN 46032 U.S.A.